

# The Similarity of Barnyardgrass Weed in Lowland Rice Fields

*by Alridiwirsa Alridiwirsa*

---

**Submission date:** 11-Oct-2023 02:15PM (UTC+0700)

**Submission ID:** 2192283612

**File name:** ss\_Weed\_in\_Lowland\_Rice\_Fields\_Based\_on\_Seed\_Characteristics.pdf (1.61M)

**Word count:** 5441

**Character count:** 24525



## The Similarity of Barnyardgrass Weed in Lowland Rice Fields Based on Seed Characteristics

Alridiwihsah<sup>(1)</sup>, Mohammad Basyuni<sup>(2,5)\*</sup>, Koko Tampubolon<sup>(3)</sup>, Novilda Elizabeth

Mustamu<sup>(4)</sup>

<sup>(1)</sup>Program Study of Agrotechnology, Faculty of Agriculture, Universitas Muhammadiyah Sumatera Utara, Medan 20238, Indonesia; <sup>(2)</sup>Program Study of Forestry, Faculty of Forestry, Universitas Sumatera Utara, Medan 20155, Indonesia; <sup>(3)</sup>Program Study of Agrotechnology, Faculty of Agriculture, Universitas Sumatera Utara, Medan 20155, Indonesia; <sup>(4)</sup>Program Study of Agrotechnology, Faculty of Science and Technology, Universitas Labuhanbatu, Rantauprapat 21415, Sumatera Utara, Indonesia; <sup>(5)</sup>Center of Excellence for Mangrove, Universitas Sumatera Utara, Medan 20155, Indonesia.



IN LOWLAND rice areas, the presence of barnyardgrass weed (*Echinochloa crus-galli*) is disadvantageous to Indonesian farmers and requires an initial assessment and identification of characteristics for management. Therefore, this study aimed to obtain the characteristics and similarities of barnyardgrass accessions from lowland rice fields based on the morphological characteristics of the seed. A descriptive-analytical method was used to determine the barnyardgrass characteristics of 45 accessions from nine sub-districts in Serdang Bedagai and one accession from Medan City (MDN), Indonesia, as a comparison. This study was conducted from May to August 2022 and the data were analyzed using one-way ANOVA, followed by Tukey's test at  $P < 0.05$ . The hierarchical cluster and constellation plot were generated using Ward's method. The results showed that 45 accessions of barnyardgrass from lowland rice fields had sizes of upper glume, lower glume, upper lemma, flag leaf length, and flag leaf width ranging from 2.93 to 4.33mm, 1.45 to 1.96mm, 0.03 to 17.10mm, 22.77 to 45.40cm, and 0.53 to 1.13cm, respectively. Among all accessions, only PG4 accession was nearest (1.825) to MDN. Furthermore, there were five clusters with 13 accessions including SB4; PC1; PC5; PG3; PG4; BK1; BK3; BK5; PB1; PB2; TB2; TB1; and TT2, classified in the same cluster as MDN. This information is very important as an initial reference for farmers in the management of barnyardgrass weeds in their lowland rice fields.

**Keywords:** Accession, Characteristics, Identification, Lowland rice, Similarity.

## Introduction

Barnyardgrass (*Echinochloa crus-galli*) is an annual weed that grows to a height of 150cm and reproduces through seeds (Bagavathiannan et al., 2012). It can grow in tropical and subtropical climates and produce up to 39,000 seeds (Bagavathiannan et al., 2013; Miller et al., 2018). Additionally, it exhibits high adaptability, rapid germination, abundant seed production, and strong

competitiveness, which can result in economic losses to the main crop (Mennan et al., 2012; Bajwa et al., 2015) and a loss of rice yield ranging from 21 to 79% (Wilson et al., 2014).

The presence of a barnyardgrass population at a distance of 40cm from lowland rice plants can decrease yield by 27%, nitrogen availability by 60 to 80%, inhibits rice growth (Wilson et al., 2014) and decrease rice yield by 30 to 100% throughout

\*Corresponding author email: m.basyuni@usu.ac.id

Received 25/09/2022; Accepted 23/01/2023

DOI: 10.21608/ejbo.2023.165026.2150

Edited by: Dr. AlBaraa ElSaied, Faculty of Science, Al-Azhar University, Cairo, Egypt.

©2023 National Information and Documentation Center (NIDOC)

the growing season (Bhullar et al., 2016; Marchesi & Saldain, 2019). *E. crus galli* var. *mitis* can lower rice grain production by 12.7 to 42.6% by inhibiting photosynthesis rate, root weight, root oxidation activity, and biomass accumulation (Zhang et al., 2017). Alridiwersah et al. (2022) added that bamyardgrass had five populations in lowland rice, with root length and volume of 39.4cm and 246.1ml and dry weight of roots and shoots of 92.7 g and 65.6g, respectively.

The presence of bamyardgrass in lowland rice fields is feared to be unfavorable to farmers in the central areas, such as Serdang Bedagai District. In 2021, Serdang Bedagai District had the second-largest rice harvest area (48,121.62ha) after Deli Serdang District from North Sumatra Province, Indonesia (Statistics of Sumatera Utara, 2022). Based on the Indonesian Center for Rice Research (2020), land preparation for lowland rice cultivation in Indonesia includes flooding the land to 2-5cm above the surface for 2-3 days before plowing the soil to a depth of 15-20cm with a tractor to kill weeds. The soil is then incubated for 3-4 days before conducting a second plowing one week after the first tilling for mudding the soil. Additionally, the ground is flattened and left moist and unflooded for 1-2 days. Rice seeds are then sown and transplanted before plant management is conducted by weed control using manual, mechanical, and chemical methods.

Herbicides are commonly used by farmers in Serdang Bedagai District to control weeds in their planting areas. Bamyardgrass is a common weed that is difficult to control. The weed must be managed appropriately using several approaches, which begin by identifying its characteristics and similarities to determine the relationship between accessions from various regions. According to Tahir (2016), the bamyardgrass from several locations in Arkansas has a leaf length of 16.2 to 30.0cm, a leaf width of 1.2-1.5 cm, upper glume of 2.94 to 3.80mm, lower glume of 1.21 to 1.68mm, an upper lemma of 2.71 to 3.55mm, and 7,186-71,494 seeds per plants. Alridiwersah et al. (2022) also added that the upper epidermis, mesophyll, lower epidermis, and stomata density of bamyardgrass leaves from lowland rice plants were 44.3  $\mu\text{m}$ , 155.6 $\mu\text{m}$ , 40.8 $\mu\text{m}$ , and 69.0n/ $\mu\text{m}$ , respectively.

A characteristics similarity approach is required <sup>31</sup> for decisions regarding weed control (Juraimi et al., 2005; Bastiani et al., 2016; Tampubolon et al.,

2019). However, the characteristics identification and similarity of bamyardgrass from lowland rice fields in Indonesia have never been reported. Therefore, an initial effort is required to determine several characters, such as flag leaf length and width, upper and lower glume, and upper lemma from various sub-districts in Serdang Bedagai. This study aimed to obtain the characteristics and similarities of bamyardgrass accessions from lowland rice fields based on the morphological characteristics of the seed. This information is expected to be a basic strategy for the management of bamyardgrass weed.

## Materials and Methods

### Seed collection area of bamyardgrass accession

The bamyardgrass accessions were obtained from nine sub-districts in Serdang Bedagai District, including Sei Bamban, Sei Rampah, Tebing Tinggi, Perbaungan, Teluk Mengkudu, Pegajahan, Pantai Cermin, Tanjung Beringin, and Bandar Khalipah, and one accession from Medan City, Indonesia, for comparison. Based on farmer interviews, the sub-district accessions were controlled with herbicide, while the comparison was controlled manually. Each sub-district consisted of five accessions from different villages, resulting in 45 accessions. Each was collected from ten populations of bamyardgrass and composited. Bamyardgrass seeds with physiological maturity criteria marked by seeds loss when <sup>28</sup>d were obtained from lowland rice fields. The flag leaf length and width of each accession were measured and environmental conditions influencing the bamyardgrass growth were recorded (Table 1). Furthermore, the bamyardgrass seeds were collected between May and August 2022 and dried for two days.

### Methodology and data analysis

A descriptive-analytical method was used to describe the similarity of bamyardgrass based on the morphological observations of the seeds, which include the upper and lower glume, upper lemma, and flag leaf length and width. Twelve seeds were randomly taken from each bamyardgrass accession, and the glume and lemma were measured using an electronic digital caliper <sup>19</sup>. The characteristics of the seeds and leaves were analyzed using one-way ANOVA (analysis of variance) followed by Tukey's test at  $P < 0.05$ . Using the JMP 17 software, the similarity of bamyardgrass weed accessions was determined using a hierarchical cluster analysis and a constellation plot with Ward's method.

TABLE 1. Locations and accession codes of barnyardgrass form Serdang Bedagai District and Medan City, Indonesia

| Sub-districts  | Accessions code | Villages            | Longitude  | Latitude  | Altitude (m asl) | Temperature (°C) | Humidity (%) | Air pressure (hPa) |
|----------------|-----------------|---------------------|------------|-----------|------------------|------------------|--------------|--------------------|
| Sei Bamban     | SB1             | Bakaran Batu        | 99°08.089' | 3°25.259' | 5                | 26.7             | 88           | 1012               |
|                | SB2             | Hapoltahan Nauli    | 99°08.383' | 3°26.552' | 4                | 25.1             | 92           | 1011               |
|                | SB3             | Gempolan            | 99°07.415' | 3°25.078' | 9                | 25.0             | 93           | 1011               |
|                | SB4             | Sei Buluh           | 99°10.605' | 3°26.848' | 2                | 25.3             | 91           | 1010               |
|                | SB5             | Sukadamai           | 99°09.636' | 3°24.762' | 6                | 24.6             | 92           | 1010               |
| Sei Rampah     | SR1             | Firdaus             | 99°08.210' | 3°30.776' | 9                | 29.1             | 68           | 1005               |
|                | SR2             | Pematang Ganjang    | 99°08.235' | 3°27.905' | 13               | 28.0             | 76           | 1006               |
|                | SR3             | Sei Parit           | 99°07.958' | 3°28.603' | 11               | 32.2             | 56           | 1006               |
|                | SR4             | Sei Rejo            | 99°09.885' | 3°30.434' | 10               | 30.7             | 65           | 1004               |
|                | SR5             | Sei Rampah          | 99°10.168' | 3°29.590' | 10               | 30.4             | 67           | 1004               |
| Tebing Tinggi  | TT1             | Paya Lombang        | 99°11.737' | 3°22.216' | 16               | 29.0             | 75           | 1009               |
|                | TT2             | Paya Mabar          | 99°12.323' | 3°23.715' | 15               | 28.9             | 75           | 1009               |
|                | TT3             | Sei Priok           | 99°13.738' | 3°24.511' | 12               | 29.9             | 67           | 1008               |
|                | TT4             | Kuta Baru           | 99°12.491' | 3°21.887' | 16               | 31.2             | 63           | 1006               |
|                | TT5             | Meriah Padang       | 99°05.111' | 3°18.138' | 30               | 29.3             | 81           | 1005               |
| Perbaungan     | PB1             | Melati I            | 98°58.034' | 3°33.567' | 10               | 24.1             | 79           | 1009               |
|                | PB2             | Melati II           | 98°58.032' | 3°33.177' | 22               | 25.1             | 79           | 1009               |
|                | PB3             | Jambur Pulau        | 98°57.755' | 3°34.322' | 1                | 25.1             | 78           | 1008               |
|                | PB4             | Cinta Air           | 98°59.606' | 3°34.488' | 16               | 26.1             | 77           | 1007               |
|                | PB5             | Citaman Jemih       | 98°57.526' | 3°33.731' | 10               | 27.1             | 63           | 1007               |
| Teluk Mengkudu | TM1             | Pasar Baru          | 99°06.396' | 3°33.056' | 9                | 30.2             | 63           | 1006               |
|                | TM2             | Pematang Gantung    | 99°07.967' | 3°33.068' | 10               | 29.9             | 64           | 1006               |
|                | TM3             | Sialang Buah        | 99°07.082' | 3°33.862' | 14               | 30.2             | 65           | 1005               |
|                | TM4             | Bongak Besar        | 99°09.038' | 3°33.060' | 17               | 29.5             | 65           | 1005               |
|                | TM5             | Sentang             | 99°08.333' | 3°33.287' | 12               | 29.5             | 65           | 1005               |
| Pegajahan      | PG1             | Bingkat IX-B        | 98°57.552' | 3°28.899' | 30               | 29.0             | 93           | 1008               |
|                | PG2             | Bingkat XI          | 98°57.512' | 3°28.546' | 13               | 28.0             | 93           | 1008               |
|                | PG3             | Bingkat X-A         | 98°57.107' | 3°28.800' | 13               | 28.0             | 93           | 1008               |
|                | PG4             | Pegajahan           | 98°56.800' | 3°28.821' | 19               | 28.0             | 93           | 1008               |
|                | PG5             | Bingkat X-B         | 98°58.427' | 3°28.729' | 26               | 28.0             | 94           | 1009               |
| Pantai Cermin  | PC1             | Kota Pari           | 98°57.625' | 3°38.446' | 14               | 30.1             | 65           | 1009               |
|                | PC2             | Pematang Kasih      | 99°01.282' | 3°37.617' | 2                | 31.2             | 67           | 1006               |
|                | PC3             | Kuala Lama          | 99°01.512' | 3°37.137' | 11               | 30.2             | 67           | 1006               |
|                | PC4             | Naga Kaisar         | 99°03.656' | 3°36.252' | 19               | 30.2             | 70           | 1007               |
|                | PC5             | Pantai Cermin Kanan | 98°58.150' | 3°38.430' | 21               | 30.1             | 71           | 1007               |

TABLE 1. Cont.

| Sub-districts    | Accessions code | Villages             | Longitude  | Latitude  | Altitude (m asl) | Temperature (°C) | Humidity (%) | Air pressure (hPa) |
|------------------|-----------------|----------------------|------------|-----------|------------------|------------------|--------------|--------------------|
| Tanjung Beringin | TB1             | Nagur                | 99°11.259' | 3°30.712' | 8                | 28.0             | 79           | 1005               |
|                  | TB2             | Mangga Dua           | 99°11.018' | 3°30.637' | 7                | 27.1             | 80           | 1006               |
|                  | TB3             | Tebing Tinggi        | 99°13.680' | 3°29.070' | 22               | 25.6             | 90           | 1009               |
|                  | TB4             | Pematang Cermai      | 99°12.650' | 3°28.709' | 6                | 25.9             | 89           | 1009               |
|                  | TB5             | Pematang Terang      | 99°14.571' | 3°26.858' | 9                | 25.8             | 90           | 1009               |
| Bandar Khalipah  | BK1             | Galam Sei Serimah    | 99°15.197' | 3°26.468' | 5                | 29.0             | 67           | 1006               |
|                  | BK2             | Pekan Bandar Kalifah | 99°16.767' | 3°26.633' | 4                | 28.9             | 68           | 1006               |
|                  | BK3             | Kayu Besar           | 99°17.133' | 3°25.996' | 1                | 29.0             | 68           | 1005               |
|                  | BK4             | Jahar                | 99°16.897' | 3°24.302' | 10               | 28.8             | 72           | 1005               |
|                  | BK5             | Bandar Tengah        | 99°15.907' | 3°23.973' | 17               | 27.4             | 81           | 1006               |
| Medan Selayang   | MDN             | Padang Bulan         | 98°39.009' | 3°33.170' | 14               | 25.0             | 34           | 1002               |

## Results and Discussion

### *Seeds morphological characters and flag leaf in barnyardgrass*

The results showed that the sizes of the glume and lemma, as well as flag leaf length and width, differed significantly between barnyardgrass accessions (Table 2). Upper and lower glume sizes were found to be the largest in TB3 and TM1 accessions (Tanjung Beringin 3 and Teluk Mengkudu 1), measuring  $4.33 \pm 0.20\text{mm}$  and  $1.96 \pm 0.02\text{mm}$ , respectively. PB1 (Perbaungan 1) had the longest upper lemma of all accessions, measuring  $17.10 \pm 0.18\text{mm}$ . Furthermore, the largest values of flag leaf length and width of barnyardgrass accession were found in PB4 and PC2, measuring  $45.40 \pm 0.66$  and  $1.13 \pm 0.03\text{cm}$ , respectively.

The difference in the size of the barnyardgrass seeds for each accession (Serdang Bedagai) with the comparison accession (MDN) can be seen visually in Fig. 1. Based on the observations of the barnyardgrass seed, the upper lemma was visible. The longest lemma from each sub-district was found in SB5; SR2; TT5; PB1; TM1; PC4; TB3; and BK2 accessions, but the upper lemma of PG1-5 did not differ from the accession comparison.

### *Correlation coefficient value*

The flag leaf length and width were positively correlated with the size of the upper glume of barnyardgrass accessions (Table 3). Similarly, both characters were significant and positively correlated ( $0.505^{**}$  and  $0.344^{*}$ ) with the size of the lower glume of barnyardgrass seeds.

### *Hierarchical cluster and similarity matrix of barnyardgrass accessions*

Based on the similarity matrix between barnyardgrass accessions from Serdang Bedagai and MDN (Table 4), only the PG4 accession was closely related (1.825). This indicates that it had the most similar characteristics to the barnyardgrass seeds from MDN. Barnyardgrass accessions were grouped into five clusters based on the hierarchical cluster analysis and constellation plot (Fig. 2). Cluster 1 had eight accessions, including SB1; SB3; SR1; SR2; TT4; PB3; TM2; and PG5, while cluster 2 consisted of five accessions including TM1; TM4; BK4; PC3; and TB5. Cluster 3 had six accessions consisting of TT1; PG1; PG2; PB4; PC2; and TB4, while Cluster 4 consisted of 13 accessions comprising TM3; TM5; SR3; SR4; SR5; TT3; TT5; SB2; SB5; PB5; BK2; TB3; and PC4. Meanwhile, cluster 5 had 14 accessions containing MDN; SB4; PC1; PC5; PG3; PG4; BK1; BK3; BK5; PB1; PB2; TB2; TB1; and TT2.

**TABLE 2.** Upper dan lower glume, upper lemma, flag leaf length, and width of barnyardgrass accessions from lowland rice fields in Serdang Bedagai District and the comparison accession (MDN), Indonesia

| Accessions | UG (mm)       | LG (mm)       | UL (mm)        | FLL (cm)        | FLW (cm)      |
|------------|---------------|---------------|----------------|-----------------|---------------|
| MDN        | 3.90±0.30 abc | 1.85±0.07 a-d | 0.04±0.01 i    | 35.57±2.56 a-e  | 0.87±0.03 ab  |
| SB1        | 3.80±0.06 abc | 1.71±0.04 a-d | 8.80±0.22 d-h  | 31.40±1.32 a-e  | 0.80±0.03 ab  |
| SB2        | 3.54±0.05 abc | 1.74±0.02 a-d | 12.94±0.74 a-d | 30.80±0.69 a-e  | 0.73±0.03 ab  |
| SB3        | 3.82±0.09 abc | 1.83±0.01 a-d | 7.20±0.24 e-h  | 31.90±1.24 a-e  | 0.83±0.03 ab  |
| SB4        | 3.84±0.08 abc | 1.76±0.03 a-d | 9.63±0.25 c-h  | 33.87±0.01 a-e  | 0.87±0.03 ab  |
| SB5        | 4.06±0.09 ab  | 1.65±0.02 a-d | 13.12±0.41 a-d | 28.23±2.89 a-e  | 0.73±0.07 ab  |
| SR1        | 4.01±0.07 abc | 1.75±0.04 a-d | 11.26±0.21 b-f | 33.53±2.49 a-e  | 0.83±0.04 ab  |
| SR2        | 3.64±0.08 abc | 1.71±0.03 a-d | 11.86±0.77 b-e | 37.27±2.05 a-e  | 0.83±0.04 ab  |
| SR3        | 3.55±0.06 abc | 1.69±0.03 a-d | 9.26±0.42 d-h  | 30.80±0.41 a-e  | 0.67±0.01 ab  |
| SR4        | 3.43±0.03 abc | 1.82±0.05 a-d | 6.12±0.07 gh   | 31.00±2.98 a-e  | 1.00±0.04 ab  |
| SR5        | 3.52±0.01 abc | 1.83±0.04 a-d | 0.03±0.00 i    | 29.57±0.46 a-e  | 0.53±0.01 b   |
| TT1        | 3.55±0.07 abc | 1.89±0.02 abc | 0.05±0.01 i    | 44.53±0.66 ab   | 1.07±0.01 a   |
| TT2        | 3.37±0.04 abc | 1.48±0.02 bcd | 5.12±0.25 h    | 39.03±0.70 a-e  | 1.03±0.01 ab  |
| TT3        | 3.52±0.07 abc | 1.46±0.02 cd  | 12.97±0.27 a-d | 29.83±0.08 a-e  | 0.67±0.01 ab  |
| TT4        | 2.95±0.09 c   | 1.57±0.02 a-d | 11.28±0.52 b-f | 30.27±2.33 a-e  | 0.83±0.10 ab  |
| TT5        | 3.40±0.09 abc | 1.55±0.02 a-d | 14.21±0.82 abc | 24.80±0.77 d-7  | 0.53±0.01 b   |
| PB1        | 3.33±0.01 abc | 1.48±0.01 bcd | 17.10±0.18 a   | 32.87±0.12 a-e  | 0.97±0.01 ab  |
| PB2        | 3.47±0.06 abc | 1.62±0.03 a-d | 13.43±0.77 a-d | 26.63±0.95 cde  | 1.03±0.04 ab  |
| PB3        | 2.93±0.08 c   | 1.57±0.04 a-d | 9.99±0.32 c-h  | 27.30±2.86 b-e  | 0.77±0.08 ab  |
| PB4        | 3.78±0.13 abc | 1.94±0.05 a   | 0.04±0.00 i    | 45.40±0.66 a    | 1.07±0.01 a   |
| PB5        | 3.62±0.10 abc | 1.90±0.04 ab  | 11.50±0.95 b-f | 39.17±0.66 a-e  | 1.00±0.03 ab  |
| TM1        | 3.09±0.13 bc  | 1.96±0.02 a   | 15.67±0.11 ab  | 31.70±0.70 a-e  | 0.80±0.03 ab  |
| TM2        | 3.18±0.06 bc  | 1.71±0.01 a-d | 11.54±0.37 b-f | 27.30±1.37 b-e  | 0.77±0.06 ab  |
| TM3        | 3.37±0.07 abc | 1.60±0.05 a-d | 6.85±0.69 fgh  | 33.60±0.57 a-e  | 1.00±0.07 ab  |
| TM4        | 3.56±0.10 abc | 1.82±0.00 a-d | 11.87±0.99 b-e | 27.70±0.68 b-e  | 0.80±0.03 ab  |
| TM5        | 2.99±0.08 bc  | 1.64±0.04 a-d | 5.81±0.36 gh   | 28.47±1.86 a-e  | 0.70±0.05 ab  |
| PG1        | 3.68±0.05 abc | 1.67±0.02 a-d | 0.07±0.01 i    | 28.83±0.77 a-e  | 0.90±0.03 ab  |
| PG2        | 3.25±0.06 abc | 1.57±0.04 a-d | 0.05±0.01 i    | 31.37±0.50 36   | 1.07±0.01 a   |
| PG3        | 3.57±0.03 abc | 1.80±0.03 a-d | 0.08±0.01 i    | 42.53±0.97 abc  | 0.93±0.01 ab  |
| PG4        | 3.51±0.07 abc | 1.78±0.03 a-d | 0.06±0.01 i    | 35.70±1.57 a-e  | 0.90±0.03 ab  |
| PG5        | 3.50±0.03 abc | 1.71±0.03 a-d | 0.04±0.01 i    | 26.73±2.83 cd-7 | 0.77±0.08 ab  |
| PC1        | 3.31±0.02 abc | 1.83±0.04 a-d | 0.06±0.01 i    | 39.13±0.85 a-e  | 0.97±0.01 ab  |
| PC2        | 3.68±0.09 abc | 1.94±0.04 a   | 0.07±0.01 i    | 40.13±0.38 a-d  | 1.13±0.03 a   |
| PC3        | 3.17±0.09 bc  | 1.55±0.04 a-d | 0.05±0.01 i    | 23.13±1.57 de-7 | 0.80±0.03 ab  |
| PC4        | 2.99±0.05 bc  | 1.47±0.02 bcd | 8.61±0.49 d-h  | 27.43±0.36 b-e  | 0.73±0.01 ab  |
| PC5        | 3.54±0.08 abc | 1.80±0.04 a-d | 0.06±0.01 i    | 24.63±1.67 de-7 | 0.87±0.03 ab  |
| TB1        | 3.29±0.07 abc | 1.58±0.04 a-d | 0.08±0.01 i    | 28.70±0.08 a-e  | 0.97±0.01 ab  |
| TB2        | 3.58±0.12 abc | 1.65±0.03 a-d | 0.04±0.01 i    | 26.10±0.49 cde  | 0.97±0.04 ab  |
| TB3        | 4.33±0.20 a   | 1.48±0.00 bcd | 12.02±0.33 b-e | 26.13±1.10 cde  | 0.67±0.04 ab  |
| TB4        | 4.02±0.10 abc | 1.67±0.05 a-d | 0.07±0.00 i    | 28.67±0.51 a-e  | 0.90±0.03 ab  |
| TB5        | 3.10±0.10 bc  | 1.68±0.05 a-d | 0.05±0.01 i    | 31.17±1.27 a-e  | 0.80±0.03 ab  |
| BK1        | 3.68±0.11 abc | 1.77±0.04 a-d | 0.06±0.01 i    | 25.40±0.52 cde  | 0.93±0.04 ab  |
| BK2        | 3.25±0.07 abc | 1.45±0.04 d   | 11.74±0.37 b-f | 22.77±0.49 e    | 0.53±0.27 1 b |
| BK3        | 4.00±0.10 abc | 1.69±0.04 a-d | 0.09±0.01 i    | 28.40±1.14 a-e  | 1.03±0.05 ab  |
| BK4        | 3.53±0.11 abc | 1.49±0.02 bcd | 10.59±0.42 c-g | 32.27±0.61 a-e  | 0.80±0.03 ab  |
| BK5        | 3.49±0.06 abc | 1.75±0.05 a-d | 0.08±0.01 i    | 28.00±1.09 b-e  | 1.03±0.03 ab  |

Note: the mean followed by different letter is significant in the Tukey's test at  $P < 0.005 \pm$  standard error. UG= upper glume; LG= lower glume; UL= upper lemma; FLL= flag leaf length, FLW= flag leaf width.




Fig. 1. The difference in the size of the seed glume and lemma in barnyardgrass accessions from Serdang Bedagai District, Indonesia and the comparison accession (MDN) visually

TABLE 3. The correlation coefficient between the barnyardgrass accessions from Serdang Bedagai District, Indonesia and the comparison accession

| Traits | FLL | FLW     | UG    | LG      | UL       |
|--------|-----|---------|-------|---------|----------|
| FLL    | 1   | 0.528** | 0.156 | 0.505** | -0.185   |
| FLW    |     | 1       | 0.110 | 0.344*  | -0.436** |
| UG     |     |         | 1     | 0.254   | -0.080   |
| LG     |     |         |       | 1       | -0.323*  |
| UL     |     |         |       |         | 1        |

Note: \*and\*\* Correlation is significant at the 0.05 and 0.01 level. FLL= flag leaf length, FLW= flag leaf width, UG= upper glume, LG= lower glume, UL= upper lemma.

TABLE 4. The similarity matrix value between the barnyardgrass accessions from Serdang Bedagai District compared to the comparison accession (MDN), Indonesia

| Accessions | Matrix value | No | Accessions | Matrix value |
|------------|--------------|----|------------|--------------|
| 1 SB1      | 4.151        | 26 | PG1        | 3.681        |
| 2 SB2      | 8.473        | 27 | PG2        | 10.584       |
| 3 SB3      | 2.123        | 28 | PG3        | 3.021        |
| 4 SB4      | 3.253        | 29 | PG4        | 1.825        |
| 5 SB5      | 10.001       | 30 | PG5        | 5.680        |
| 6 SR1      | 4.549        | 31 | PC1        | 4.419        |
| 7 SR2      | 5.956        | 32 | PC2        | 4.627        |
| 8 SR3      | 7.589        | 33 | PC3        | 15.327       |
| 9 SR4      | 4.833        | 34 | PC4        | 20.879       |
| 10 SR5     | 7.879        | 35 | PC5        | 5.469        |
| 11 TT1     | 5.818        | 36 | TB1        | 9.449        |
| 12 TT2     | 12.000       | 37 | TB2        | 6.506        |
| 13 TT3     | 16.903       | 38 | TB3        | 17.747       |
| 14 TT4     | 17.800       | 39 | TB4        | 3.410        |
| 15 TT5     | 22.047       | 40 | TB5        | 8.795        |
| 16 PB1     | 19.415       | 41 | BK1        | 4.456        |
| 17 PB2     | 13.647       | 42 | BK2        | 27.019       |
| 18 PB3     | 19.117       | 43 | BK3        | 4.261        |
| 19 PB4     | 5.600        | 44 | BK4        | 11.736       |
| 20 PB5     | 5.988        | 45 | BK5        | 5.280        |
| 21 TM1     | 15.179       |    |            |              |
| 22 TM2     | 12.882       |    |            |              |
| 23 TM3     | 8.225        |    |            |              |
| 24 TM4     | 7.651        |    |            |              |
| 25 TM5     | 14.563       |    |            |              |




Fig. 2. Hierarchical clustering and constellation plot of 46 barnyardgrass accessions from Serdang Bedagai District and Medan City (comparison), Indonesia [Codes of barnyardgrass accessions and their localities are listed in Table 1]

The results showed that the sizes of the upper and lower glume, upper lemma, and flag leaf length and width of the barnyardgrass accession from Serdang Bedagai District ranged from 2.93 to 4.33mm, 1.45 to 1.96mm, 0.03 to 17.10mm, 22.77 to 45.40cm, and 0.53 to 1.13cm, respectively (Table 2). The upper and lower glume sizes in TB3 and TM1, as well as the upper lemma found in PB1 accessions, were larger than in other accessions. The results also revealed that PB4 and PC2 accessions had the highest flag leaf length and width values when compared to other accessions. This indicates that the size of seed glume from these locations could be due to the total assimilating productivity of the large flag leaf length and width. This finding was supported by the flag leaf length and width, which are positively correlated (0.156 and 0.110) with the size of the upper glume, and positively correlated and significant (0.505\*\* and 0.344\*) with the size of the lower glume (Table 3). Asana (1968) supported this result by stating that flag leaves play an important role in providing photosynthesis products to seeds. According to Ashrafuzzaman et

al. (2009) and Rahman et al. (2013), flag leaf area was positively correlated with panicle length and seed product. In contrast, the size of the upper lemma was negatively correlated with the flag leaf length and width (Table 3). It was assumed that the photosynthate product from the flag leaf influenced glume formation and panicle length. Liu et al. (2021) also reported that flag leaf length and width were positively correlated to panicle length in *Echinochloa* ecotypes.

It was observed that only the PG4 accession was closely related to MDN, with a similarity value of 1.825. The hierarchical cluster confirmed these findings, with the SB4; PC1; PC5; PG3; PG4; BK1; BK3; BK5; PB1; PB2; TB2; TB1; and TT2 accessions clustered in one group with the MDN. This indicates that the characters of the upper and lower glume, upper lemma, flag leaf length, and width of the 13 accessions were not significantly different from MDN (Table 2). This was consistent with a study by Tahir (2016) which stated that eight of 95 *Echinochloa sp* accessions from several locations in Arkansas were clustered

with the barnyardgrass based on vegetative and generative characteristics. Altop & Mennan (2011) classified barnyardgrass weeds from lowland rice fields in Turkey into three clusters based on morphological characters, with 28 of 34 barnyardgrass populations having similarities. The findings also showed that the morphological character approach is almost similar when using a molecular approach (RAPD), with 33 of 34 barnyardgrass populations clustered into one group. Liu et al. (37) also characterized the morphophysiology of *E. crus-galli*; *E. colona*; and *E. muricata* from rice-growing in Texas, producing five clusters and discovered that the *E. crus-galli* (ECH22) ecotype was related to *E. muricata* (ECH33). Claerhout et al. (2016) identified three clusters of 14 barnyardgrass accessions based on morphological characters from maize fields in Belgium.

Based on the result, 13 of 45 accessions from lowland rice fields in Serdang Bedagai District were similar to MDN. This result showed that farmers' management of barnyardgrass weeds from these accessions did not differ significantly to the comparison accession. These results are reported for the first time in Indonesia. However, the morphological character approach to assessing similarity should be re-examined using molecular methods.

### Conclusions

Based on morphological characters, 45 accessions of barnyardgrass weeds from Serdang Bedagai District had different sizes of upper and lower glume, upper lemma, and flag leaf length and width ranged from 2.93-4.33mm; 1.45-1.96mm; 0.03-17.10mm; 22.77-45.40cm; and 0.53-1.13cm, respectively. Accessions of TB3; TM1; PB1; PB4; and PC2 had the largest characteristic size compared to other accessions. Additionally, the PG4 accession was closely related to MDN (1.825). It was also found that the SB4; PC1; PC5; PG3; PG4; BK1; BK3; BK5; PB1; PB2; TB2; TB1; and TT2 accessions were classified into one cluster with MDN.

**4**  
**21** **nowledgments:** The authors would like to thank the Ministry of Education, Culture, Research, and Technology, (Indonesia) for funding this study with the 4mber 153/E5/PG.02.00.PT/2022. In addition, the authors would also like to thank Fransisca Natalia Sihombing, S.Pd., M.P., Josefin

Batara Raja Siagian, Emil Salim Hasibuan, S.Pt., and Febri Ardiansyah Harahap, S.Pt. who have assisted in the field.

**Competing interests:** The authors report no conflicts of interest regarding this work.

**Authors' contributions:** Conceptualization and methodology (Koko Tampubolon, Alridiwersah, Mohammad Basyuni). Climate collecting and measuring (Novilda Elizabeth Mustamu, Alridiwersah) Seed collecting and measuring (Mohammad Basyuni, Koko Tampubolon) Validation and data analysis (Koko Tampubolon, Novilda Elizabeth Mustamu). Writing, review, editing, and prepare manuscript for publishing (all authors).

**Ethics approval:** Not applicable.

### References

- Alridiwersah., Tampubolon, K., Zulkifi, T.B.H., Risnawati., Yusuf, M. (2022) Allelopathic effects of *Mikania micrantha* Kunth on barnyardgrass and lowland rice. *Pesquisa Agropecuária Tropical*, **52**, e71356. <https://revistas.ufg.br/pat/article/view/71356>.
- 18**  
Altóp, E.K., Mennan, H. (2011) Genetic and morphologic diversity of *Echinochloa crus-galli* populations from different origins. *Phytoparasitica*, **39**(1), 93-102.
- 25**  
Asana, R.D. (1968) In quest of yield. *Indian Journal of Plant Physiology*, **11**(1&2), 1-9.
- 11**  
Ashrafuzzaman, M., Islam, M.R., Ismail, M.R., Shahidullah, S.M., Hanafi, M.M. (2009) Evaluation of six aromatic rice varieties for yield and yield contributing characters. *International Journal of Agriculture and Biology*, **11**(5), 616-620.
- 8**  
Bagavathiannan, M.V., Norsworthy, J.K., Smith, K.L., Neve, P. (2012) Seed production of barnyardgrass (*Echinochloa crus-galli*) in response to time of emergence in cotton and rice. *The Journal of Agricultural Science*, **150**(6), 717-724.
- 10**  
Bajwa, A.A., Jabran, K., Shahid, M., Ali, H.H., Chauhan, B.S. (2015) Eco-biology and management of *Echinochloa crus-galli*. *Crop Protection*, **75**, 151-162.
- 15**  
Bastiani, M.O., Lamego, F.P., Agostinetto, D., Langaro,

- A.C., da Silva, D.C. (2016) Relative competitiveness of soybean cultivars with barnyardgrass. *Bragantia*, **75**(4), 435-445.
- 13 Bhullar, M.S., Kumar, S., Kaur, S., Kaur, T., Singh, J., Yadav, R., Chauhan, B.S., Gill, G. (2016) Management of complex weed flora in dry-seeded rice. *Crop Protection*, **83**, 20-26.
- 12 Claerhout, S., Dewaele, K., De Riek, J., Reheul, D., De Cauwer, B. (2016) Morphological and genetic variability of local *Echinochloa* accessions and the link with herbicide sensitivity. *Weed Research*, **56**(2), 137-148.
- Indonesian Center for Rice Research. (2020). Recommendations for rice cultivation in various agroecosystems. Ministry of Agriculture, Subang, West Java, Indonesia. 62p.
- 9 Juraimi, A.S., Tasrif, A., Kadir, J., Sastroutomo, S.S., Napis, S. (2005) Morphological and RAPD variability among Malaysian ecotypes of barnyardgrass (*Echinochloa crus-galli* var. *crus-galli* (L.) P. Beauv.). *Plant Protection Quarterly*, **20**(2), 52-57.
- 1 Liu, R., Singh, V., Abugho, S., Lin, H. S., Zhou, X. G., Bagavathiannan, M. (2021) Morphophysiological diversity and its association with herbicide resistance in *Echinochloa* ecotypes. *Weed Science*, **70**(1), 26-35.
- 22 Marchesi, C., Saldain, N.E. (2019) First report of herbicide-resistant *Echinochloa crus-galli* in Paraguayan rice fields. *Agronomy*, **9**(12), 790. <https://doi.org/10.3390/agronomy9120790>.
- Mennan, H., Ngouajio, M., Sahin, M., Isik, D., Altop, E.K. (2012) Competitiveness of rice (*Oryza sativa* L.) cultivars against *Echinochloa crus-galli* (L.) Beauv. in water-seeded production systems. *Crop Protection*, **41**, 1-9. <https://doi.org/10.1016/j.cropro.2012.04.027>.
- 17 Miller, M.R., Norsworthy, J.K., Scott, R.C. (2018) Evaluation of florypyrauxifen-benzyl on herbicide-resistant and herbicide-susceptible barnyardgrass accessions. *Weed Technology*, **32**(2), 126-134.
- 6 Rahman, M.A., Haque, M.E., Sikdar, B., Islam, M.A., Matin, M.N. (2013) Correlation analysis of flag leaf with yield in several rice cultivars. *Journal of Life and Earth Science*, **8**, 49-54.
- Statistics of Sumatera Utara. (2022) Sumatera Utara Province in figures 2022. Medan, Indonesia.
- 1 Tahir, H. (2016) Characterization of *Echinochloa* spp. in Arkansas. Thesis. University of Arkansas.
- 24 Tampubolon, K., Sihombing, F.N., Siburian, E., Sulastri, Y.S., Purba, Z., Samosir, S.T.S., Karim, S. (2019) Similarity and phylogenetic analysis of herbicide-resistant goosegrass (*Eleusine indica*) biotypes. *Caraka Tani: Journal of Sustainable Agriculture*, **34**(2), 162-169.
- 14 Wilson, M.J., Norsworthy, J.K., Scott, R.C., Gbur, E.E. (2014) Program approaches to control herbicide-resistant barnyardgrass (*Echinochloa crus-galli*) in midsouthern United States rice. *Weed Technology*, **28**(1), 39-46.
- Zhang, Z., Gu, T., Zhao, B., Yang, X., Peng, Q., Li, Y., Bai, L. (2017) Effects of common *Echinochloa* varieties on grain yield and grain quality of rice. *Field Crops Research*, **203**, 163-172.

# The Similarity of Barnyardgrass Weed in Lowland Rice Fields

## ORIGINALITY REPORT

15%  
SIMILARITY INDEX

%  
INTERNET SOURCES

15%  
PUBLICATIONS

%  
STUDENT PAPERS

## PRIMARY SOURCES

- 1 Rui Liu, Vijay Singh, Seth Abugho, Hao-Sheng Lin, Xin-Gen Zhou, Muthukumar Bagavathiannan. " Morphophysiological diversity and its association with herbicide resistance in ecotypes ", Weed Science, 2021  
Publication 1%
- 2 Martina Huber, Magdalena M. Julkowska, L. Basten Snoek, Hans van Veen et al. "Towards increased shading capacity: A combined phenotypic and genetic analysis of rice shoot architecture", PLANTS, PEOPLE, PLANET, 2023  
Publication 1%
- 3 Wan Arfiani Barus, Koko Tampubolon, Alridiwirsa Alridiwirsa, Tengku Boumedine Hamid Zulkifli et al. "Seedlings and Vegetative Stage of Red Rice as Affected by Induction Gamma-Rays under Saline Stress", Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis, 2023  
Publication 1%

4

Novilda Elizabeth Mustamu, Koko Tampubolon, Alridiwersah, Mohammad Basyuni et al. "Drought stress induced by polyethylene glycol (PEG) in local maize at the early seedling stage", Heliyon, 2023

Publication

1 %

5

Tao Song, Feiyun Xu, Wei Yuan, Moxian Chen, Qijuan Hu, Yuan Tian, Jianhua Zhang, Weifeng Xu. "Combining alternate wetting and drying irrigation with reduced phosphorus fertilizer application reduces water use and promotes phosphorus use efficiency without yield loss in rice plants", Agricultural Water Management, 2019

Publication

1 %

6

Babyrani Panda, Subhankar Mondal, Ashish Mohanty, Jeetendra Senapaty et al. "Exploring the physiological basis of yield enhancement in New Generation Rice (NGR): a comparative assessment with non-NGR rice genotypes", Plant Physiology Reports, 2023

Publication

1 %

7

Juan J. Rios, Agatha Agudelo, Diego A. Moreno, Micaela Carvajal. "Growing broccoli under salinity: the influence of cultivar and season on glucosinolates content", Scientia Agricola, 2020

Publication

1 %

8

Aurora M. Baltazar, Surajit K. Datta. "Biology and Management of Some Problem Weeds", Wiley, 2023

Publication

1 %

9

Emine Kaya Altop, Husrev Mennan. "Genetic and morphologic diversity of *Echinochloa crus-galli* populations from different origins", *Phytoparasitica*, 2010

Publication

1 %

10

Hengzhi Wang, Yi Li, Lipeng Wang, Weitang Liu, Jinxin Wang. "Baseline sensitivity of *Echinochloa crus-galli* (L.) P.Beauv. to tripyrasulfone, a new HPPD-inhibiting herbicide, in China", *Crop Protection*, 2022

Publication

1 %

11

A Roy, MAR Sarkar, SK Paul. "Effect of age of seedlings at staggered transplanting and nutrient management on yield performance of aromatic fine rice (cv. BRRI dhan38)", *SAARC Journal of Agriculture*, 2018

Publication

1 %

12

Ivan Hoste, Filip Verloove. "Taxonomy of the weed species of the genus *Echinochloa* (Poaceae, Paniceae) in Southwestern Europe: Exploring the confused current state of affairs", *PhytoKeys*, 2022

Publication

1 %

13 SANJEEV KUMAR, SHIVANI SHIVANI, J S MISHRA, SANTOSH KUMAR, UJJWAL KUMAR, R C BHARATI. "Efficacy of pre- and post-emergence herbicides on complex weed flora in direct-seeded rice (*Oryza sativa*) in the eastern plains", The Indian Journal of Agricultural Sciences, 2018

Publication

---

14 Babita Patni, Sudhir Kumar Guru, Marcello Iriti, Sara Vitalini. "Elicitation of the Allelopathic Potential of Rice by Methyl Salicylate Treatment", Applied Sciences, 2019

Publication

---

15 Leandro Galon, Carlos Orestes Santin, André Luiz Radünz, André Andres et al. "Competitive ability of sweet sorghum (*Sorghum bicolor* (L.) Moench) cultivars against hairy beggarticks (*Bidens pilosa* L.)", Revista de Ciencias Agrícolas, 2022

Publication

---

16 Xinxin Tang, Rong Gong, Wenqiang Sun, Chaopu Zhang, Sibin Yu. "Genetic dissection and validation of candidate genes for flag leaf size in rice (*Oryza sativa* L.)", Theoretical and Applied Genetics, 2017

Publication

---

17 Hannah E. Wright, Jason K. Norsworthy, Trenton L. Roberts, Robert C. Scott, Jarrod T.

Hardke, Edward E. Gbur. "Use of florpyrauxifen-benzyl in non-flooded rice production systems", Crop, Forage & Turfgrass Management, 2021

Publication

---

18

Shuhong He, Hongbin Lang, Jiatao Shen, Luqing Zhang, Hua Fang, Yunlong Yu. "Herbicidal activity of atrazine to barnyard grass depends upon soil characteristics", Pest Management Science, 2022

Publication

---

19

Basyuni, Mohammad, Shigeyuki Baba, Yuji Kinjo, Lollie A.P. Putri, Luthfi Hakim, and Hirosuke Oku. "Salt-dependent increase in triterpenoids is reversible upon transfer to fresh water in mangrove plants Kandelia candel and Bruguiera gymnorhiza", Journal of Plant Physiology, 2012.

Publication

---

20

Rahman, Md Asadur, ME Haque, B Sikdar, Md Asadul Islam, and Muhammad Nurul Matin. "Correlation Analysis of Flag Leaf with Yield in Several Rice Cultivars", Journal of Life and Earth Science, 2014.

Publication

---

21

A A Oksari, D Susanty, F H Rizki, I F Wanda, Arinana, Dadang. "Potential of Dioscorea bulbifera L. as a bio-insecticide in controlling

<1 %

<1 %

<1 %

<1 %

dry wood termites (*Cryptotermes cynocephalus* Ligh.)", IOP Conference Series: Earth and Environmental Science, 2023

Publication

---

22

Rana Nadeem Abbas, Asif Iqbal, Muhammad Aamir Iqbal, Omar M. Ali, Raees Ahmed, Raina Ijaz, Adel Hadifa, Brandon J. Bethune. "Weed-Free Durations and Fertilization Regimes Boost Nutrient Uptake and Paddy Yield of Direct-Seeded Fine Rice (*Oryza sativa* L.)", *Agronomy*, 2021

Publication

---

23

S F Batubara, F Agus, A Rauf, D Elfiati. "Soil respiration and microbial population in tropical peat under oil palm plantation", IOP Conference Series: Earth and Environmental Science, 2019

Publication

---

24

Alridiwirsah, Koko Tampubolon, Novilda Elizabeth Mustamu, Mujiyo, Mehdizadeh Mohammad. "Weed diversity as affected by tillage and ammonium glufosinate herbicide", *Pesquisa Agropecuária Tropical*, 2022

Publication

---

25

IF Wardlaw. "Carbohydrate Storage and Mobilisation by the Culm of Wheat Between Heading and Grain Maturity: the Relation to Sucrose Synthase and Sucrose-Phosphate

<1 %

<1 %

<1 %

<1 %

## Synthase", Australian Journal of Plant Physiology, 1994

Publication

26

Dilshan Benaragama, Brian G. Rossnagel, Steven J. Shirtliffe. "Breeding for Competitive and High-Yielding Crop Cultivars", Crop Science, 2014

Publication

<1 %

27

E.A.E. Elsheikh, A.H. El Tinay, I.A. Fadul. "Effect of nutritional status of faba bean on proximate composition, anti-nutritional factors and in vitro protein digestibility (IVPD)", Food Chemistry, 1999

Publication

<1 %

28

Elena R. Sanchez Olguin. "Assessment of gene flow from a herbicide-resistant indica rice (*Oryza sativa* L.) to the Costa Rican weedy rice (*Oryza sativa*) in Tropical America: factors affecting hybridization rates and characterization of F1 hybrids", Transgenic Research, 03/29/2009

Publication

<1 %

29

Pierre, Johann S., Jai Perroux, Alex Whan, Anne L. Rae, and Graham D. Bonnett. "Poor Fertility, Short Longevity, and Low Abundance in the Soil Seed Bank Limit Volunteer Sugarcane from Seed", Frontiers in Bioengineering and Biotechnology, 2015.

Publication

<1 %

---

|    |                                                                                                                                                                                                                   |      |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 30 | Qiongnan Gu, Shihai Chu, Qichao Huang, Anan Chen, Lin Li, Ruhai Li. "Colletotrichum Echinochloae: A Potential Bioherbicide Agent for Control of Barnyardgrass (Echinochloa Crus-Galli (L.) Beauv.)", Plants, 2023 | <1 % |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|

---

Publication

|    |                                                                                                                                                                                                                                                                                                                                                              |      |
|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 31 | Zhou, Da-Yong, Dong-Dong Ma, Jun Zhao, Xiu-Lin Wan, Lei Tong, Shuang Song, Jing-Feng Yang, and Bei-Wei Zhu. "Simultaneous Recovery of Protein and Polysaccharide from Abalone (Haliotis Discus Hannai Ino) Gonad Using Enzymatic Hydrolysis Method : Recovery of Valuable Components from Abalone Gonad", Journal of Food Processing and Preservation, 2015. | <1 % |
|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|

---

Publication

|    |                                                                            |      |
|----|----------------------------------------------------------------------------|------|
| 32 | "Rice Production Worldwide", Springer Science and Business Media LLC, 2017 | <1 % |
|----|----------------------------------------------------------------------------|------|

---

Publication

|    |                                                                                                                                                      |      |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| 33 | E. Santen. "Genetic Diversity in Tall Fescue Ecotypes from Alabama I. Maturity, Morphological Traits, and Disease Reaction", Plant Breeding, 11/1991 | <1 % |
|----|------------------------------------------------------------------------------------------------------------------------------------------------------|------|

---

Publication

|    |                                                                                      |      |
|----|--------------------------------------------------------------------------------------|------|
| 34 | Emel Yıldız, Hasan Ayyıldız. "chapter 14 Electronic Word-of-Mouth", IGI Global, 2014 | <1 % |
|----|--------------------------------------------------------------------------------------|------|

---

Publication

---

35

Lu Wang. "Mapping 49 quantitative trait loci at high resolution through sequencing-based genotyping of rice recombinant inbred lines", Theoretical and Applied Genetics, 09/28/2010

Publication

<1 %

36

Pil Nam Seong, Kyoung Mi Park, Geun Ho Kang, Soo Hyun Cho, Beom Young Park, Hyun Seok Chae, Hoa Van Ba. "The Differences in Chemical Composition, Physical Quality Traits and Nutritional Values of Horse Meat as Affected by Various Retail Cut Types", Asian-Australasian Journal of Animal Sciences, 2015

Publication

<1 %

37

Spencer C. H. Barrett, Blake F. Wilson. "Colonizing ability in the - complex (barnyard grass). I. Variation in life history ", Canadian Journal of Botany, 1981

Publication

<1 %

38

Yusuff Oladosu, M. Y. Rafii, Norhani Abdullah, Mohammad Abdul Malek et al. "Genetic Variability and Selection Criteria in Rice Mutant Lines as Revealed by Quantitative Traits", The Scientific World Journal, 2014

Publication

<1 %

39

Moses M. Muraya, Hartwig H. Geiger, Fabrice Sagnard, Lassana Toure et al. "Adaptive values of wild×cultivated sorghum (Sorghum bicolor (L.) Moench) hybrids in generations

<1 %

# F1, F2, and F3", Genetic Resources and Crop Evolution, 2011

Publication

---

---

Exclude quotes      Off

Exclude matches      Off

Exclude bibliography      Off