

Agronomic Character of Ratoon Rice_Stem Cutting Sizes and Seprint Liquid Organic Fertilizer

by Alridiwirsah Alridiwirsah

Submission date: 11-Oct-2023 02:49PM (UTC+0700)

Submission ID: 2192296582

File name: ice_Stem_Cutting_Sizes_and_Seprint_Liquid_Organic_Fertilizer.pdf (1.11M)

Word count: 7351

Character count: 35611

Article ID: 747
DOI: 10.5586/aa.747

Publication History
Received: 2020-04-29
Accepted: 2021-02-13
Published: 2021-06-04

Handling Editor
Zuzanna Parzymies; University of Life Sciences in Lublin, Poland; <https://orcid.org/0000-0002-4835-3804>

Authors' Contributions
A: coordinating the research, designing ideas and methods, conceptualization; KT: conceptualization, analysis and interpretation of data, field coordinator, writing, reviewing, and editing of the manuscript; FNS: data collection and analysis, writing and reviewing of the manuscript; AAS and ZP: data collection

Funding
The author would like to thank the University of Muhammadiyah Sumatera Utara for funding this research.

Competing Interests
No competing interests have been declared.

Copyright Notice
The Author(s) 2021. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits redistribution, commercial and noncommercial, provided that the article is properly cited.

ORIGINAL RESEARCH PAPER in PLANT STRUCTURE

Agronomic Character of Ratoon Rice: Stem Cutting Sizes and Seprint Liquid Organic Fertilizer

Alridiwersah^{1*}, Koko Tampubolon²,
Fransisca Natalia Sihombing³, Andi Agus Suprianto¹,
Zavandri Purba⁴

¹ Program Study of Agrotechnology, Faculty of Agriculture, University of Muhammadiyah Sumatera Utara, Medan 20238, Sumatera Utara, Indonesia

² Program Study of Agrotechnology, Faculty of Agriculture and Animal Husbandry, Universitas Jut Nyak Dhien, Medan 20123, Sumatera Utara, Indonesia

³ Master's Program of Agribusiness, Faculty of Agriculture, Universitas Sumatera Utara, Medan 20155, Sumatera Utara, Indonesia

⁴ Program Study of Agrotechnology, Faculty of Agriculture, Universitas Sumatera Utara, Medan 20155, Sumatera Utara, Indonesia

* To whom correspondence should be addressed. Email: alridiwersah@umsu.ac.id

Abstract

This study aimed to identify the optimum rice stem cutting size combined with the application of Seprint liquid organic fertilizer (SLOF) to promote the growth and yield of ratoon rice. This research was conducted in a greenhouse at the Faculty of Agriculture, University of Muhammadiyah Sumatera Utara, North Sumatra Province, Indonesia, from January to August 2018. The experiment was set up as a completely randomized design factorial with four replications. The first factor consisted of the cutting sizes of the rice stem (R1 = 5 cm, R2 = 10 cm, and R3 = 15 cm) and the second factor involved the SLOF dosage (S1 = 5 mL L⁻¹, S2 = 10 mL L⁻¹, and S3 = 15 mL L⁻¹). The parameters were analyzed using analysis of variance (ANOVA), and the means were determined by Duncan's multiple range test (DMRT). The results showed that the R2 treatment significantly increased the ratoon rice height at 10 to 30 days after cutting (DAC), the number of tillers at 40 to 60 DAC, the number of productive tillers, and the yield per hectare. The S2 and S3 applications significantly increased the yield per hill and ratoon rice height at 20 DAC. The R2S1 and R2S2 combination treatments significantly increased the number of filled grains per panicle and the ratoon rice height at 10 DAC. Cutting the stems at 10 cm combined with various dosages of SLOF resulted in greater growth and yield of ratoon rice for achieving an adequate food supply.

Keywords

dosage; growth; yield; crop model; optimization technique; ratoon technology

1. Introduction

Food is a basic human need, and adequate, nutritious, and safe food can lead to high-quality human resources. Grain (rice) is a major source of energy. According to United States Department of Agriculture (USDA) data, countries with the highest level of rice consumption worldwide from 2015 to 2019 are located in Asia, specifically in China (29.66%), India (26.39%), Indonesia (7.88%), Bangladesh (7.32%), and Vietnam (4.54%) (Ministry of Agriculture of the Republic of Indonesia, 2019). In Indonesia, rice is the main food requirement for more than 90% of the population. The consumption of rice per capita has decreased from 107.71 kg capita⁻¹ year⁻¹ in 2002 to 96.33 kg capita⁻¹ year⁻¹ in 2018 (Ministry of Agriculture of the Republic of Indonesia, 2019). However, domestic rice yield continues to increase, although it is trending toward a slower growth rate. Indonesia's population

growth increased linearly from 2005 to 2015 (226.71 to 258.16 million), and the projected population until 2035 is estimated to be up to 305.65 million (Statistics Indonesia, 2019). The total domestic consumption of Indonesian rice continues to increase, even though the per capita consumption shows a decrease.

The increased domestic consumption of rice in Indonesia must be followed by a program to increase rice yield via intensification. One of the intensification programs that can be followed is system cultivation of ratoon rice. Mengel and Wilson (1981) stated that ratoon is a technique of harvesting production from tillers from the main plant stumps to increase grain yield without increasing land area. The ratoon technique has been used to shorten the age of the rice harvest. The maturing of hybrid rice varieties is required at 105–124 days after scattering or approximately 4 months (Indonesian Center for Rice Research, 2019). Therefore, in a year, there can only be two lots of rice planting until harvesting. The ratoon system can produce three to four lots of rice harvesting. The advantages of the ratoon system are the higher efficiency of resource use per unit area of land and unit per timing (Santos et al., 2003), reduction in land preparation, cost savings on planting seeds (Zandstra & Samson, 1979), and time saving of planting because of decreased growth time (Haque & Coffman, 1980). In addition, ratoon rice is preferred by farmers because of its high quality, including cooking and flavor, short growth period, reduced water and fertilizer costs, and decreased pest control (Yazdpour et al., 2012). Alizadeh and Habibi (2016) reported that the grain thickness and amylose content of ratoon rice were higher, ranging from 1.18% to 2.25% and 2.28% to 7.69%, respectively, compared to the main crop.

System cultivation of ratoon rice has been reported. Pratama et al. (2018) stated that the time of cutting the remaining stumps on day 10 of the rice harvest was significant for obtaining the largest root diameter of 0.89 mm with a stump cutting height of 4 cm above the soil surface and could increase grain productivity by 3.39 to 3.54 tons ha^{-1} . Nuzul et al. (2018) stated that the highest yield components (e.g., the number and length of panicles, harvest age, grain number per panicle, filled grain, 1,000-grains weight, total dry weight, and harvest index), and rice yield (grain per hectare) were achieved in the cutting at harvest of a height 3 cm above the soil surface with a grain yield of 3.54 tons ha^{-1} . Susilawati et al. (2012) stated that the cutting height of ratoon rice was related to the internode of the plant, and the nodes contained lateral shoots that produced ratoon shoots. Jichao and Xiaohui (1996) showed that the number of growing shoots was determined by the cutting height stems of the main plant; however, the condition was strongly influenced by residual assimilates as reserves in the stem that could be utilized for ratoon growth, and the vigor rate of the ratoon. The stem cutting height of the main plant can also affect the number of tillers and seed yield (Harrell et al., 2009) and trigger dormant shoot growth (Mahadevappa & Yogeesh, 1988).

However, in these studies, the rice cultivation system was not separated from fertilizer use to achieve optimal yield. Islam et al. (2008) reported that fertilizer application of $\text{N} = 130 \text{ kg ha}^{-1}$, $\text{P}_2\text{O}_5 = 75 \text{ kg ha}^{-1}$, $\text{K}_2\text{O} = 80 \text{ kg ha}^{-1}$, $\text{S} = 13 \text{ kg ha}^{-1}$, and $\text{Zn} = 4 \text{ kg ha}^{-1}$ significantly increased the yield of Boro rice ratoon by 1.42 tons ha^{-1} . The high price of fertilizer with limited availability has resulted in the use of liquid organic fertilizer as an alternative to increase the availability and efficiency of irrigation rice nutrient uptake. The effect of liquid organic fertilizer on the growth and yield of irrigated rice has been widely reported (Ginting, 2019; Mahmoodi et al., 2020; Niis & Nik, 2017; Oh & Kim, 2013; Solihin et al., 2019); however, studies regarding the use of liquid organic fertilizer on the growth and yield of ratoon rice are limited. Thus, a special assessment of the influence of specific liquid organic fertilizers is required to increase the growth and yield of ratoon rice. The present study aimed to identify the optimum size of the rice stem cutting (ratoon) combined with the application of Seprint liquid organic fertilizer to promote growth and yield of ratoon in rice variety Inpari 30.

2. Material and Methods

2.1. Experiment I (Preliminary Research Stage)

The main plant included in the present study was the lowland rice variety Inpari 30. This variety was used because it has an average yield of up to 7.2 tons ha^{-1} (Indonesian Agency for Agricultural Research and Development, 2020), which is higher than that of the Ciherang, Situ Bagendit, and Pepe varieties (Pramono et al., 2011). In addition, the taste of this rice variety is preferred by farmers in the Cicantayan subdistrict compared to the Inpari 22, Inpari 24, and Inpari 25 varieties (Putra & Haryati, 2018) from the Assessment Institute for Agricultural Technology Pasar Miring, Pagar Merbau subdistrict, Deli Serdang District, North Sumatra Province, Indonesia. The main plant was placed in a 20-L water bucket with an upper diameter of 31 cm, a lower diameter of 22 cm, and a height of 31 cm (Figure 1A). This experiment was conducted in a greenhouse at the Faculty of Agriculture, University of Muhammadiyah Sumatera Utara, Indonesia, from January to August 2018. The bucket was filled with soil up to 5 cm from the surface. Two lowland rice seedlings were planted in the bucket and inundated with water until the vegetative phase of the main plant. The water in the bucket was then decreased to the bearing capacity for harvesting. The main plant was harvested 115 days after planting, followed by cutting the rice stem (Experiment II).




Figure 1 Size of the bucket (A), design of the second experiment (B), and Seprint liquid organic fertilizer by Bunga Tani (C).

2.2. Experiment II (Main Research Stage)

After the main rice was harvested, it was cut from the rice stem base, as shown in Figure 1B. Seprint liquid organic fertilizer (SLOF) was applied 1 week after cutting the rice stems by dissolving in 1 L of water and pouring onto the soil surface. The nutrient content of SLOF was 11% N, 2% K_2O , 0.17% P_2O_5 , 0.49% Bioret, 8.98% boron (B), 5.35% cobalt (Co), 5.83% manganese (Mn), <0.01 ppm molybdenum (Mo), 18.16 ppm zinc (Zn), and 9.94 copper (Cu) (Figure 1C) (source: Bunga Tani). The bucket was inundated when cutting the rice stems until the end of the vegetative phase, and then the water was decreased to harvest the ratoon rice.

2.3. Research Design and Data Analysis

The present study was performed in a completely randomized design factorial with 15 replications. The first factor consisted of the cutting sizes of the rice stem ($\text{R}_1 = 5 \text{ cm}$, $\text{R}_2 = 10 \text{ cm}$, and $\text{R}_3 = 15 \text{ cm}$), and the second factor involved the SLOF dosage ($\text{S}_1 = 5 \text{ mL L}^{-1}$, $\text{S}_2 = 10 \text{ mL L}^{-1}$, and $\text{S}_3 = 15 \text{ mL L}^{-1}$). The distance between the treatments was $50 \times 50 \text{ cm}$. The parameters measured included the height and growth rate of the ratoon rice; the number of tillers at 10 to 60 days after cutting (DAC); the yield component of ratoon rice including the number of productive tillers, number of grains per panicle, number of filled grains per panicle, number of empty grains per panicle, 1,000-grains weight, yield per hill, and yield per hectare. The agronomic traits of ratoon rice are transformed in the form of natural logarithms (ln) due to the different units and then analyzed with the Pearson

correlation. The calculation of the growth rate and yield per hectare of the ratoon rice can be seen in Equation (1) and Equation (2):

$$\text{Growth rate (cm day}^{-1}\text{)} = \frac{\text{Average height of ratoon rice}}{\text{Days of cutting}} \quad (1)$$

$$\text{Yield (ha}^{-1}\text{)} = NP \times NPT \times NFGP \times 1,000 \text{ GW} \quad (2)$$

Notes:

- NP – number population per hectare = $\frac{\text{Land area (ha}^{-1}\text{)}}{\text{Plant spacing}} = \frac{10,000 \text{ m}^2}{0.5 \text{ m} \times 0.5 \text{ m}} = 40,000$ population;
- NPT – number of productive tillers;
- $NFGP$ – number of filled grains per panicle;
- 1,000 GW – 1,000-grains weight (g).

The parameters were analyzed using ANOVA, and the mean was determined by DMRT at 5% significance, and correlation analysis for agronomic traits of ratoon rice using IBM SPSS statistics v.20 software.

3. Results

3.1. Height and Growth Rate of Ratoon Rice

Based on the ANOVA results, the cutting of the stems significantly increased the height of the ratoon rice at 10 to 30 DAC. In contrast, no significant increase was found at 40 and 60 DAC. SLOF application significantly increased the height of the ratoon rice at 20 DAC. In contrast, no significant increase was found at 10, 30, 40, and 60 DAC. The interaction of the cutting of the rice stems with SLOF application significantly increased the height of the ratoon rice at 10 DAC. However, no significant increase was found at 20 to 60 DAC (Table 1). The R2 treatment had the highest ratoon rice height at 10 to 60 DAC compared to the other cutting sizes, with significant increases from 10 to 30 DAC. The S3 application had the highest ratoon rice height at 10 to 40 DAC compared to the other doses, although it was significantly increased only at 20 DAC. The highest ratoon rice height was found in the S2 application at 60 DAC. The R2S2 and R2S3 combination treatments significantly increased the ratoon rice height to 27.98 cm and 25.55 cm, respectively, at 10 DAC. The highest ratoon rice heights were found in the R2S3 treatment at 20 to 60 DAC.

3.2. Number of Tillers of Ratoon Rice

Based on the ANOVA results, the cutting of the rice stems significantly increased the number of tillers of ratoon rice at 40 and 60 DAC. In contrast, no significant increase was found at 10 to 30 DAC. The SLOF application and interaction with stem cutting did not significantly increase the number of tillers of ratoon rice (Table 2). The average number of tillers of lowland rice before becoming a ratoon ranged from 9.25 to 11.75 tillers. The R2 treatment had the highest number of tillers of ratoon rice at 10 to 60 DAC compared to the other cutting sizes, with significant increases at 40 to 60 DAC. The S2 application had the highest number of tillers of ratoon rice (25.33 tillers) at 60 DAC compared to the other doses. The R2S3 combination treatment had the highest number of tillers of ratoon rice at 36.25 tillers at 60 DAC compared to the other combinations.

3.3. Yield Component of Ratoon Rice

Based on the ANOVA results, the cutting of the rice stems significantly affected the number of productive tillers. In contrast, the SLOF dosages and interactions with stem cuttings did not significantly increase the number of productive tillers of ratoon rice (Figure 2–Figure 4). The R2 treatment had the highest number of productive tillers (24.92 tillers) compared to the other rice stem cuttings.

Based on the ANOVA results, the cutting of the rice stems and SLOF dosages did not significantly affect the number of filled grains per panicle. In contrast, their interactions significantly increased the number of filled grains per panicle of ratoon

Table 1 Effect of rice stem cutting and Seprint liquid organic fertilizer (SLOF) dosages on the ratoon rice height from 10 to 60 days after cutting (DAC).

Treatment	Ratoon rice height (cm)			Average	Growth rate (cm day ⁻¹)
	S1	S2	S3		
10 days after cutting					
R1	9.46 d–h	7.05 h	10.33 c–h	8.95 b	0.89
R2	7.45 gh	27.98 a	25.55 a	20.33 a	2.03
R3	13.13 b–h	8.90 fgh	9.45 e–h	10.49 b	1.05
Average	10.01 ns	14.64 ns	15.11 ns		
20 days after cutting					
R1	15.53 ns	17.35 ns	22.93 ns	18.60 b	0.93
R2	15.25 ns	42.35ns	48.80 ns	35.47 a	1.77
R3	27.90 ns	12.73 ns	35.40 ns	25.34 ab	1.27
Average	19.56 b	24.14 ab	35.71 a		
30 days after cutting					
R1	31.28 ns	39.95 ns	36.88 ns	36.04 b	1.20
R2	39.70 ns	58.33 ns	63.63 ns	53.89 a	1.80
R3	46.40 ns	38.13 ns	49.93 ns	44.82 ab	1.49
Average	39.13 ns	45.47 ns	50.15 ns		
40 days after cutting					
R1	46.90 ns	52.98 ns	50.73 ns	50.20 ns	1.26
R2	61.90 ns	63.93 ns	75.23 ns	67.02 ns	1.68
R3	60.70 ns	54.28 ns	55.83 ns	56.94 ns	1.42
Average	56.50 ns	57.06 ns	60.60 ns		
60 days after cutting					
R1	59.20 ns	65.80 ns	58.18 ns	61.06 ns	1.02
R2	67.50 ns	70.38 ns	77.53 ns	71.80 ns	1.20
R3	67.93 ns	66.28 ns	64.63 ns	66.28 ns	1.10
Average	64.88 ns	67.49 ns	66.78 ns		

Note: Data followed by different letters in the same column and row are significantly different according to the DMRT at 5% significance. Size cutting of rice stems (R1 = 5 cm, R2 = 10 cm, and R3 = 15 cm). SLOF dosages (S1 = 5 mL L⁻¹, S2 = 10 mL L⁻¹, and S3 = 15 mL L⁻¹). ns = not significant.

rice (Figure 2–Figure 4). The R2S1 combination treatment significantly increased the highest number of filled grains per panicle to 114.30 grains compared to the other combinations. The R2 treatment had the highest number of filled grains per panicle (97.58 grains) compared to the other cutting sizes. The S2 application had the highest number of filled grains per panicle (91.53 grains) compared to the other doses.

Based on the ANOVA results, the SLOF dosages significantly increased the yield per hill of ratoon rice. In contrast, the cutting of the rice stems and their interactions did not significantly increase the yield per hill of ratoon rice (Figure 2–Figure 4). The S2 application significantly had the highest yield per hill of ratoon rice of 54.32 g compared to the other doses. The R2 treatment had the highest yield per hill of ratoon rice of 46.56 g compared to the other cutting sizes. The R2S2 combination treatment had the highest yield per hill of ratoon rice of 60.63 g compared to the other combinations.

Based on the ANOVA results, the cutting of rice stems significantly increased the yield per hectare of ratoon rice. In contrast, the SLOF dosages and their interactions did not significantly increase the yield per hectare of ratoon rice (Figure 5). The R2 treatment had the highest yield per hectare of ratoon rice of 3.58 tons ha⁻¹ compared to the other cutting sizes. The S2 application had the highest yield per hectare of ratoon rice of 3.12 tons ha⁻¹ compared to the other dosages, whereas the R2S3 combination treatment had the highest yield per hectare of ratoon rice of 3.88 tons ha⁻¹ compared to the other combinations.

Table 2 Number of tillers of ratoon rice after cutting the stems and Seprint liquid organic fertilizer (SLOF) dosages from 10 to 60 days after cutting (DAC).

Treatment	Number of tillers of ratoon rice			Average
	S1	S2	S3	
10 days after cutting				
R1	6.75 ns	2.00 ns	3.75 ns	4.17 ns
R2	7.00 ns	7.00 ns	6.00 ns	6.67 ns
R3	8.00 ns	5.00 ns	4.50 ns	5.83 ns
Average	7.25 ns	4.67 ns	4.75 ns	
20 days after cutting				
R1	8.75 ns	4.25 ns	5.00 ns	6.00 ns
R2	8.25 ns	14.50 ns	15.25 ns	12.67 ns
R3	15.00 ns	9.00 ns	8.75 ns	10.92 ns
Average	10.67 ns	9.25 ns	9.67 ns	
30 days after cutting				
R1	9.00 ns	7.25 ns	8.00 ns	8.08 ns
R2	9.00 ns	19.25 ns	24.00 ns	17.50 ns
R3	20.00 ns	12.75 ns	15.00 ns	15.92 ns
Average	12.75 ns	13.08 ns	15.67 ns	
40 days after cutting				
R1	10.75 ns	12.00 ns	10.50 ns	11.08 b
R2	10.50 ns	33.25 ns	33.00 ns	25.58 a
R3	22.75 ns	23.00 ns	19.50 ns	21.75 ab
Average	14.67 ns	22.75 ns	21.00 ns	
60 days after cutting				
R1	13.00 ns	13.25 ns	13.25 ns	13.17 b
R2	16.00 ns	35.00 ns	36.25 ns	29.08 a
R3	24.00 ns	27.75 ns	23.50 ns	25.08 a
Average	17.67 ns	25.33 ns	24.33 ns	

Note: Data followed by different letters in the same column and row are significantly different according to the DMRT at 5%. Size cutting of rice stems (R1 = 5 cm, R2 = 10 cm, and R3 = 15 cm). SLOF dosages (S1 = 5 mL L⁻¹, S2 = 10 mL L⁻¹, and S3 = 15 mL L⁻¹). ns = not significant.

Based on the ANOVA results, the stem cutting sizes, SLOF dosages, and their interactions were insignificant on the yield components of ratoon rice (number of grains per panicle, number of empty grains per panicle, and 1,000-grains weight; Figure 2–Figure 4). The R2 treatment had the highest number of grains per panicle and 1,000-grains weight of 116.77 grains and 25.91 g, respectively, compared to the other cutting sizes. The R3 treatment had the highest number of empty grains per panicle (20.29 grains) compared to the other cutting sizes. The S1 application had the highest number of grains per panicle and number of empty grains per panicle of 113.59 and 22.69 grains, respectively, compared to the other doses. The S2 application had the highest 1,000-grains weight of 25.92 g compared to the other doses. The R2S1 combination treatment had the highest number of grains per panicle of 139.85 grains compared to the other combinations. The R3S1 and R3S2 combination treatments had the highest number of empty grains per panicle and 1,000-grains weight of 25.90 grains and 26.78 g, respectively, compared to the other combinations.

3.4. Correlations of Agronomic Traits of Ratoon Rice

The agronomic traits of ratoon rice include the plant height, number of tillers per panicle, number of productive tillers, number of grains per panicle, number of filled grains per panicle, 1,000-grains weight, and yield per hill significantly increase the yield per hectare of ratoon rice with the highest contribution was found in the number of




Figure 2 Effect of rice stem cutting sizes on the yield component of ratoon rice: (A) number of productive tillers; (B) number of grains per panicle; (C) number of filled grains per panicle; (D) number of empty grains per panicle; (E) 1,000-grains weight; (F) yield per hill. Data followed by different letters on the same graph are significantly different according to DMRT at 5% $\pm$ standard error (SE).




Figure 3 Effect of Seprint liquid organic fertilizer (SLOF) dosages on the yield component of ratoon rice: (A) number of productive tillers; (B) number of grains per panicle; (C) number of filled grains per panicle; (D) number of empty grains per panicle; (E) 1,000-grains weight; (F) yield per hill. Data followed by different letters on the same graph are significantly different according to DMRT at 5% $\pm$ standard error (SE).

productive tillers (correlation coefficient value of 0.885) (Table 3). The present study confirmed that an increase the yield per hectare of ratoon rice along with an increase in the number of productive tillers in the combination treatment of stem cutting with SLOF dosages.




Figure 4 Combinations of stem cutting sizes and Seprint liquid organic fertilizer (SLOF) dosages on the yield component of ratoon rice: (A) number of productive tillers; (B) number of grains per panicle; (C) number of filled grains per panicle; (D) number of empty grains per panicle; (E) 1,000-grains weight; (F) yield per hill. Data followed by different letters on the same graph are significantly different according to DMRT at 5% ± standard error (SE).




Figure 5 Effect of rice stem cutting sizes on the yield per hectare of ratoon rice. Data followed by different letters on the same line are significantly different according to DMRT at 5% ± standard error (SE).

Table 3 Correlations of agronomic traits of ratoon rice in the treatment of stems cutting and Seprint liquid organic fertilizer (SLOF) dosages.

Agronomic traits of ratoon rice	PH	NT	NPT	NGP	NFGP	NEGP	1,000 GW	YH	Y
PH	1								
NT	0.500**	1							
NPT	0.683**	0.612**	1						
NGP	-0.043	-0.143	0.157	1					
NFGP	0.104	-0.051	0.217	0.927**	1				
NEGP	-0.264	-0.207	0.055	0.693**	0.387*	1			
1,000 GW	0.167	0.243	0.271	-0.085	0.090	-0.311	1		
YH	0.208	0.241	0.222	0.071	0.258	-0.310	0.329*	1	
Y	0.582**	0.500**	0.885**	0.488**	0.600**	0.137	0.396*	0.360*	1

Note: Correlation is significant at the * 0.05 and ** 0.01 level (two-tailed). Agronomic traits: PH – plant height; NT – number of tillers; NPT – number of productive tillers; NGP – number of grains per panicle; NFGP – number of filled grains per panicle; NEGP – number of empty grains per panicle; 1,000 GW – 1,000-grains weight; YH – yield per hill; Y – yield per hectare.

4. Discussion

Cutting 10 cm of the rice stem (R2 treatment) significantly increased the height of ratoon rice at 10 to 30 DAC and significantly increased the number of tillers of ratoon rice at 40 to 60 DAC and the number of productive tillers. The results were similar for the growth rate of ratoon rice, with the R2 treatment higher by 1.20 to 2.03 cm day⁻¹ compared to the R1 and R3 treatments that ranged from 0.89 to 1.26 cm day⁻¹ and 1.05 to 1.49 cm day⁻¹, respectively. This was caused by an optimal carbohydrate supply from the emergence of 30 ratoon shoots that resulted in higher vegetative growth and was supported by the plant height, number of tillers, growth rate, and number of productive tillers on the R2 treatment being higher than the other cutting sizes (Table 1, Table 2 and Figure 2–Figure 4). The optimal cutting size of rice stems supports the growth and yield of ratoon rice. Santos et al. (2003) showed a decrease in yield ranging from 12% to 37% in <5 cm rice stem cuttings, whereas cuttings >20 cm of the rice stem produced tillers faster, and the number of grains was less than the optimal cutting size. Harrell et al. (2009) found that height cutting of the main plant stems affected the number of tillers and seed yield. Susilawati et al. (2012) stated that the height and number of productive tillers for ratoon rice in genotype IPB106-F-8-1 at a 10 cm cutting was higher than that at 20 cm and 30 cm cuttings. Mareza et al. (2016) stated that cutting 40 to 50 cm of the main rice stem was significant for the number of productive tillers, with the highest found at the 40 cm cutting (6.55 tillers) compared to the other cutting sizes.

The R2 treatment significantly increased the yield per hectare of ratoon rice to 3.58 tons ha⁻¹ compared to the other cutting sizes (Figure 5). The yield per hectare of ratoon rice in the study was smaller than that of the main plants (rice variety Inpari 30) where the initial planting was 5.08 tons ha⁻¹ and the yield decreased by 29.53% compared to the main plant. This resulted in a growth pattern (height and number of tillers) of ratoon rice that was higher in the R2 treatment compared to the other cutting sizes because photosynthesis is caused the formation of assimilates and the production of filled grains. The number of productive tillers, number of filled grains per panicle, and 1,000-grains weight for the R2 treatment were also higher than the other cutting sizes (Figure 2–Figure 4). According to Harrell et al. (2009), the height of the cutting stems of the main plant affects the number of tillers and grains.

Susilawati et al. (2012) showed that the yield per hectare of ratoon rice in genotype IPB106-F-8-1 for the 10 cm cutting was greater (2.1 tons ha⁻¹) than that for the 20 and 30 cm cuttings (1.6 and 1.1 tons ha⁻¹, respectively). Nuzul et al. (2018) found that the yield of ratoon rice decreased by 58.72% from that of the main plants.

The S3 treatment significantly increased the height of ratoon rice at 20 DAC (Table 1). SLOF has a main nutrient content of 11% N, which is higher than that of

K (2%). Therefore, a SLOF dose of 15 mL L⁻¹ could support the photosynthesis level and increase the vegetative phase of the ratoon rice. According to Surtinah (2009), liquid organic fertilizers are very efficient because plants easily absorb such fertilizers. Parman (2007) stated that liquid organic fertilizers improved soil physical, chemical, and biological properties, and increased the yield and quality of plants. Niis and Nik (2017) showed that the application of liquid organic fertilizer at a dose of 6 L ha⁻¹ significantly increased the highest number of seeds per panicle and the dry seed weight of rice by 107.38 seeds and 29.65 g, respectively, compared to doses of 2 and 4 L ha⁻¹.

The S2 application significantly increased the yield per hill of ratoon rice by 54.32 g. SLOF contains 11% N and 2% K, which played an important role in improving the nutrient uptake and soil organic matter and increasing the availability of macro- and micronutrients for ratoon rice. Liquid organic fertilizers can play a role in cell division and increase assimilation, leading to increased biomass and yield of ratoon rice. According to Vidigal et al. (2002), the growth of onions increased progressively with an increased application of K fertilizer. El-Bassiony (2006) also stated that the application of K fertilizer increased the dry weight of onions. Martínez-Alcántara et al. (2016) stated that liquid organic fertilizers increased the uptake of macro- and micronutrients, and organic matter in soil compared to chemical fertilizers. Phibunwatthanawong and Riddech (2019) showed that liquid organic fertilizers contained micro- and supplementing nutrients, beneficial microorganisms such as plant growth-promoting rhizobacteria, and hormone indole acetic acid, which are not found in chemical fertilizers.

The R2S2 combination treatment significantly increased the ratoon rice height to 27.98 cm at 10 DAC. The R2S2 and R2S3 combination treatments had the highest ratoon rice height compared to the other combinations. This was caused by the main nutrients N and K in the S2 and S3 applications promoting the photosynthesis rate to form assimilates to stimulate ratoon rice growth, including the plant height, number of tillers, and number of productive tillers (Table 1, Table 2 and Figure 2–Figure 4). According to Nurhayati (2009), liquid organic fertilizer can promote metabolism in plants via the nutrient solubility process, and nutrient transport will occur from the roots to all parts of the plant. Hamzah (2014) stated that liquid organic fertilizer could increase plant height, and the number of branches and soybean pods.

The R2S1 combination treatment significantly increased the number of filled grains per panicle by 114.30 grains. This was because SLOF stimulated the metabolic process with the growth rate of ratoon rice, and photosynthesis formed assimilates and led to grain panicle production. According to Susilawati et al. (2012), the cutting of 10 cm of rice stem in genotype IPB106-F-8-1 significantly increased the number of grain per panicle, which was 108.0 grains higher than the 20 cm and 30 cm cuttings (78.5 and 71.1 grains, respectively). Marengga et al. (2016) stated that cutting 10 cm of the rice stems significantly improved the panicle length and the number of grains per panicle of ratoon rice by 14.93 cm and 47.05 grains, respectively, compared to the cuttings of 20 to 50 cm.

The correlations result confirmed that the number of productive tillers had a positive linear correlation (0.885) and was classified as very strong in increasing the yield per hectare of ratoon rice compared to another agronomic traits (Table 3). Similar results were reported by Sadeghi (2011); Ranawake et al. (2013) the number of productive tillers and the number of grains per panicle had a significant effect on grain yield per plant. Faruq et al. (2014) also reported that the number of productive tillers per hill has a correlated positively and significant increase the grain yield per plant of ratoon rice.

The application of ratoon rice technology combined with SLOF greatly encouraged rice yield to improve adequate food supply for a year of rice cultivation (main plants and ratoon technology) can produce an average yield of 8.66 tons ha⁻¹. Thus, farmers will receive a double benefit by applying ratoon technology during a cycle of irrigated rice.

5. Conclusions

The R2 treatment significantly increased the ratoon rice height from 20.33 to 53.89 cm at 10 to 30 DAC, the number of tillers from 25.58 to 29.08 tillers at 40 to 60 DAC, the number of productive tillers to 24.92 tillers, and the yield per hectare of ratoon rice to 3.58 tons ha⁻¹. The S3 application significantly increased the ratoon rice height to 35.71 cm at 20 DAC, and the S2 application significantly increased the yield per hill of ratoon rice to 54.32 g. The R2S2 combination treatment significantly increased the ratoon rice height to 27.98 cm at 10 DAC and the R2S1 combination treatment also significantly increased the number of filled grains per panicles of ratoon rice to 114.30 grains. The R2 treatment combined with various dosages of SLOF resulted in higher growth and yield of ratoon rice compared to the other cutting sizes.

References

- Alizadeh, M. R., & Habibi, F. (2016). A comparative study on the quality of the main and ratoon rice crops. *Journal of Food Quality*, 39(6), 669–674. <https://doi.org/10.1111/jfq.12250>
- El-Bassiony, A. M. (2006). Effect of potassium fertilization on growth, yield and quality of onion plants. *Journal of Applied Sciences Research*, 2(10), 780–785.
- Faruq, G., Taha, R. M., & Prodhan, Z. H. (2014). Rice ratoon crop: A sustainable rice production system for tropical hill agriculture. *Sustainability*, 6(9), 5785–5800. <https://doi.org/10.3390/su6095785>
- Ginting, J. (2019). Effect of liquid organic fertilizer on the rice varieties field production. *Conference Series: Earth and Environmental Science*, 260, Article 012177. <https://doi.org/10.1088/1755-1315/260/1/012177>
- Hamzah, S. (2014). Liquid organic fertilizer and manure chicken influence on the growth and yield of soybean (*Glycine max* L.). *Agrium*, 18(3), 228–234.
- Haque, M. M., & Coffman, W. R. (1980). Varietal variation and evaluation procedures for ratooning ability in rice. *SABRAO Journal*, 12(2), 113–120.
- Harrell, D. L., Bond, J. A., & Blanche, S. (2009). Evaluation of main-crop stubble height on ratoon rice growth and development. *Field Crops Research*, 114(3), 396–403. <https://doi.org/10.1016/j.fcr.2009.09.011>
- Indonesian Agency for Agricultural Research and Development. (2020). *Inpari 30 Ciherang Sub 1*. <https://www.litbang.pertanian.go.id/varietas/848/>
- Indonesian Center for Rice Research. (2019). *Klasifikasi umur tanaman padi* [Age classification of rice plants]. <http://bbpadi.litbang.pertanian.go.id/index.php/info-berita/tahukah-anda/820/klasifikasi-umur-tanaman-padi>
- Islam, M. S., Hasanuzzaman, M., & Rokonuzzaman, M. (2008). Ratoon rice response to different fertilizer doses in irrigated condition. *Agriculturae Conspectus Scientificus*, 73(4), 197–202.
- Jichao, Y., & Xiaohui, S. (1996). Effect of cutting node and leaves retained on the mother stem on rice ratooning. *Journal of Sichuan Agricultural University*, 4(7), 42–53.
- Mahadevappa, M., & Yogeesh, H. S. (1996). Rice ratooning breeding, agronomic practice, and seed production potential. In W. H. Smith, V. Kumble, & E. P. Cervantes (Eds.), *Rice ratooning* (pp. 177–186). IRRRI.
- Mahmoodi, B., Moballeghi, M., Eftekhari, A., & Neshai-Mogadam, M. (2020). Effects of foliar application of liquid fertilizer on agronomical and physiological traits of rice (*Oryza sativa* L.). *Acta Agrobotanica*, 73(3), Article 7332. <https://doi.org/10.5586/aa.7332>
- Mareza, E., Djafar, Z. R., Suwignyo, R. A., & Wijaya, A. (2016). Rice ratoon yield response to main crops cutting height in tidal swamp using direct seeding system. *AGRIVITA, Journal of Agricultural Science*, 38(2), 126–132. <https://doi.org/10.17503/agrivitya.v38i2.502>
- Martínez-Alcántara, B., Martínez-Cuenca, M. R., Bermejo, A., Legaz, F., & Quinones, A. (2016). Liquid organic fertilizers for sustainable agriculture: Nutrient uptake of organic versus mineral fertilizers in citrus trees. *PLoS One*, 11(10), Article e0161619. <https://doi.org/10.1371/journal.pone.0161619>
- Mengel, D. B., & Wilson, F. E. (1981). Water management and nitrogen fertilization of ratoon crop rice. *Agronomy Journal*, 73(6), 1008–1010. <https://doi.org/10.2391/08900314-19817361008>
- Ministry of Agriculture of the Republic of Indonesia. (2019). *Buletin konsumsi* [Food consumption bulletin]. <http://epublikasi.setjen.pertanian.go.id/arsip-buletin/53-buletin-konsumsi/620-buletin-konsumsi-vol-10-no-1-2019>
- Niis, A., & Nik, N. (2017). Pengaruh dosis dan frekuensi aplikasi pupuk organik cair (POC) terhadap pertumbuhan dan hasil padi (*Oryza sativa*, L.) [Effect of dosage and

- frequency of liquid organic fertilizer (LOF) application on growth and yield of rice (*Oryza sativa* L.]. *Savana Cendana*, 2(1), 4–7. <https://doi.org/10.32938/sc.v2i01.77>
- Nurhayati. (2009). Pengaruh cekaman air pada jenis tanah terhadap pertumbuhan dan hasil kedelai [*Glycine max* (L.) Merrill] [Effect of water stress at two types of soil on growth and yield of soybean, *Glycine max* (L.) Merrill]. *Jurnal Floratek*, 4(1), 55–64.
- Nuzul, V. S., Indradewa, D., & Kastono, I. (2018). Pengaruh waktu dan tinggi pemotongan tunggul terhadap komponen hasil dan hasil padi (*Oryza sativa* L.) Ratoon [Effects of stem cutting time and height on yield components and yield of rice ratoon system (*Oryza sativa* L.)]. *Vegetalika*, 7(2), 54–65. <https://doi.org/10.22146/veg.35773>
- Oh, T., & Kim, C. (2013). Effect of using organic fertilizer on the growth of rice and soil. *Korean Journal of Crop Science*, 58(1), 36–42. <https://doi.org/10.7740/kjcs.2013.58.1.036>
- Parman, S. (2007). Pengaruh pemberian pupuk organik terhadap pertumbuhan dan produksi kentang (*Solanum tuberosum* L.) [The effect of liquid organic fertilizer on growth and yield of potatoes (*Solanum tuberosum* L.)]. *Buletin Anatomi dan Fisiologi*, 15(2), 21–31.
- Phibunwatthanawong, T., & Riddech, N. (2019). Liquid organic fertilizer production for growing vegetables under hydroponic condition. *International Journal of Recycling of Organic Waste in Agriculture*, 8(4), 369–380. <https://doi.org/10.1007/s40093-019-0257-7>
- Pramono, J., Norma, M., & Abadi. (2011). Increased productivity of lowland rice through the introduction of new superior varieties and improved farm management in Kendal Regency. In *Prosiding Seminar Nasional: Pemberdayaan Petani Melalui Inovasi Teknologi Spesifik Lokasi* [Proceedings of the National Seminar: Empowering farmers through location-specific technology innovations] (pp. 106–113). Institute for Agricultural Technology (BPTP) in Yogyakarta; College of Agricultural Extension (STPP) in Magelang.
- Pratama, A. B., Indradewa, D., & Ambarwati, E. (2018). Karakter morfologi akar dan hasil padi raton (*Oryza sativa* L.) pada perbedaan waktu dan tinggi pemotongan tunggul sisa panen [The effect of cutting time and height of rice stubble on root morphology and yield of rice ratoon (*Oryza sativa* L.)]. *Vegetalika*, 7(4), 12–25. <https://doi.org/10.22146/veg.41150>
- Putra, S., & Haryati, Y. (2018). Kajian produktivitas dan respon petani terhadap padi varietas unggul baru di Kecamatan Parungkuda dan Cicantayan Kabupaten Sukabumi [Research on the productivity and farmers' response towards new high-yield varieties of rice in Sukabumi Regency, West Java Province]. *Jurnal Pengkajian dan Pengembangan Teknologi Pertanian*, 21(1), 1–10. <https://doi.org/10.21082/jppt.v21n1.2018.p1-10>
- Ranawake, A., Amarasingha, U., & Dahanayake, N. (2013). Agronomic characters of some traditional rice (*Oryza sativa* L.) cultivars in Sri Lanka. *Journal of the University of Ruhuna*, 1(1), 3–9. <https://doi.org/10.4038/jur.v1i1.6150>
- Sadeghi, S. M. (2011). Heritability, phenotypic correlation and path coefficient studies for some agronomic characters in landrace rice varieties. *World Applied Sciences Journal*, 13(5), 1229–1233.
- Santos, A. B., Fageria, N. K., & Prabhu, A. S. (2003). Rice ratooning management practices for higher yields. *Communications in Soil Science and Plant Analysis*, 34(5–6), 881–918. <https://doi.org/10.1081/CSS-120018981>
- Solihin, E., Yuniarti, A., Damayani, M., & Rosniawaty, S. (2019). Application of liquid organic fertilizer and N, P, K to properties of soil chemicals and growth of rice plant. *IOP Conference Series: Earth and Environmental Science*, 393, Article 012026. <https://doi.org/10.1088/1755-1315/393/1/012026>
- Statistics Indonesia. (2019). *Indonesian population projections*. Retrieved May 25, 2021, from <https://www.bps.go.id/statictable/2014/02/18/1274/proyeksi-penduduk-menurut-provinsi-2010-2035.html>
- Surtinah. (2009). Pemberian pupuk organik Super Natural Nutrition (SNN) pada tanaman selada (*Lactuca sativa* L.) di tanah Ultisol [Application of Super Natural Nutrition (SNN) organic fertilizer on lettuce (*Lactuca sativa* L.) in Ultisol soil]. *Jurnal Ilmiah Pertanian*, 6(1), 20–25.
- Susilawati, S., Purwoko, B. S., Aswiddinnoor, H., & Santosa, E. (2012). Rice ratoon production affected by stem cutting height at harvest. *Indonesian Journal of Agronomy*, 40(1), 1–7.
- Vidigal, S. M., Pereira, P. R. G., & Pacheco, D. D. (2002). Mineral nutrition and fertilization of onion. *Informe Agropecuaria*, 23(218), 36–50.
- Yazdpour, H., Shahri, M. M., Soleymani, A., & Shahrajabian, M. H. (2012). Effects of harvesting time and harvesting height on grain yield and agronomical characters in rice ratoon (*Oryza sativa* L.). *Journal of Food, Agriculture and Environment*, 10(1), 438–440.
- Zandstra, H. G., & Samson, B. T. S. (1979). Rice ratoon management. In *International Rice Research Conference, April 1979* (pp. 16–21). International Rice Research Institute.

Agronomic Character of Ratoon Rice_Stem Cutting Sizes and Seprint Liquid Organic Fertilizer

ORIGINALITY REPORT

19%
SIMILARITY INDEX

%
INTERNET SOURCES

19%
PUBLICATIONS

%
STUDENT PAPERS

PRIMARY SOURCES

- 1 Caleb Vanlalrinngama, Banshidhar Jha, S. K. Singh, A. Tigga, Bishawajit Kumar, Namata Kumari, M. K. Singh. "Trait association studies in diverse genotypes of rice for their utilization in biofortification", Environment Conservation Journal, 2023
Publication 1%
- 2 Marguerita Leavitt, Beatriz Moreno-García, Colby W. Reavis, Michele L. Reba, Benjamin R.K. Runkle. "The effect of water management and ratoon rice cropping on methane emissions and yield in Arkansas", Agriculture, Ecosystems & Environment, 2023
Publication 1%
- 3 Catarina Esteves, David Fangueiro, Mariana Mota, Miguel Martins, Ricardo P. Braga, Henrique Ribeiro. "Partial replacement of chemical fertilizers with animal manures in an apple orchard: Effects on crop performance and soil fertility", Scientia Horticulturae, 2023
Publication 1%

4

Iswanto Iswanto, Nurhayati Nurhayati, Rahmad Setia Budi. "Pengaruh Pemberian Air dan Beberapa Jenis Pupuk Organik terhadap Pertumbuhan dan Produksi Tanaman Kedelai (*Glycine max* L. Merril)", Daun: Jurnal Ilmiah Pertanian dan Kehutanan, 2021

Publication

1 %

5

Domenic T. Sanchez, Larry B. Peconcillo Jr, John V. De Vera. "Effects of Natural and Home-Made Bio-Inoculants on Containerized Okra Plants (*Abelmoschus Esculentus*)", Current Agriculture Research Journal, 2022

Publication

1 %

6

Giovani O Silva, Antonio C Bortoletto, Agnaldo DF Carvalho, Arione S Pereira. "Effect of potassium sources on potato tuber yield and chip quality", Horticultura Brasileira, 2018

Publication

<1 %

7

W. Han. "Chlorophyll a fluorescence responses of *Haloxylon ammodendron* seedlings subjected to progressive saline stress in the Tarim desert highway ecological shelterbelt", Photosynthetica, 12/2010

Publication

<1 %

8

F. A. Bahar, S. K. De Datta. " Prospects of Increasing Tropical Rice Production through Ratooning ", Agronomy Journal, 1977

Publication

<1 %

9

Acep Atma Wijaya, Haris Maulana, Gatut Wahyu Anggoro Susanto, Dadang Sumardi, Suseno Amien, Dedi Ruswandi, Agung Karuniawan. "Grain yield stability of black soybean lines across three agroecosystems in West Java, Indonesia", Open Agriculture, 2022

Publication

<1 %

10

Eduardo Martins da Silva Kaue, Crestani Mota Marcelo, Bonadeu Francismeire, Galvao de Souza Karina et al. "Effect of selective micronutrients on productivity of upland rice varieties", African Journal of Agricultural Research, 2022

Publication

<1 %

11

S. Fukai, L. Li, P. T. Vizmonte, K. S. Fischer. "Control of Grain Yield by Sink Capacity and Assimilate Supply in Various Rice () cultivars", Experimental Agriculture, 2008

Publication

<1 %

12

Kaifu Song, Guangbin Zhang, Jing Ma, Shaobing Peng, Shihua Lv, Hua Xu. "Greenhouse gas emissions from ratoon rice fields among different varieties", Field Crops Research, 2022

Publication

<1 %

13

Nadezhda Golubkina, Carmine Amalfitano, Agnieszka Sekara, Alessio Tallarita et al. "Yield and bulb quality of storage onion cultivars as

<1 %

affected by farming system and nitrogen dose", Scientia Horticulturae, 2021

Publication

14

Yue-chao WANG, Xiu-fen LI, TARPLEY Lee, Shao-bing PENG, Fu-gen DOU. "Effects of nitrogen management on the ratoon crop yield and head rice yield in South USA", Journal of Integrative Agriculture, 2021

Publication

15

Mukherjee, Bhaskar. "LiBe-14: A novel microdosimeter using LiF and BeO thermoluminescence dosimeter pairs for clinical and aerospace applications", Radiation Measurements, 2015.

Publication

16

Mapegau Mapegau, Asrizal Paiman, Marlina Marlina. "KARAKTER AGRONOMI PADI RATUN (ORIZA SATIVA. L) VARIETAS LOKAL PASANG SURUT TANJUNG JABUNG TIMUR", Jurnal Agroecotania : Publikasi Nasional Ilmu Budidaya Pertanian, 2018

Publication

17

O. Salem, R. Ghodia. "EFFECT OF FOLIAR SPRAY ON ONION PRODUCTIVITY UNDER RAINFALL AND SUPPLEMENTARY IRRIGATION CONDITIONS", Journal of Plant Production, 2012

Publication

<1 %

<1 %

<1 %

<1 %

18	A Suriadi, F Zulhaedar, M Nazam, A Hipi. "Optimal irrigation at various soil types for soybean production", IOP Conference Series: Earth and Environmental Science, 2021 Publication	<1 %
19	David B. Jones. "Rice Ratoon Response to Main Crop Harvest Cutting Height", Agronomy Journal, 1993 Publication	<1 %
20	Hanoi National University of Education Publication	<1 %
21	Shrestha, S., N. M. M. Thin, and P. Deb. "Assessment of climate change impacts on irrigation water requirement and rice yield for Ngamoeyeik Irrigation Project in Myanmar", Journal of Water and Climate Change, 2014. Publication	<1 %
22	Maria Delviani Tou. "Faktor-Faktor yang Mempengaruhi Produksi Usahatani Padi Sawah di Desa Angkaes Kecamatan Weliman Kabupaten Malaka", AGRIMOR, 2017 Publication	<1 %
23	Pepi Nur Susilawati, Resmayeti Purba. "Combination of Organic Fertilizer, Anorganic Fertilizer and Bio-Fertilizer in Maize Cultivation in Dry Land, Banten", E3S Web of Conferences, 2021 Publication	<1 %

24

S. Singh, R. S. Sengar. " Evaluation of Selected Indian Bread Wheat (L.) Genotypes for Morpho-physiological and Biochemical Characterization under Salt Stress Conditions ", Cereal Research Communications, 2016

Publication

<1 %

25

A Nurhuda, A Wibowo, Kustiyo. "Spatial-temporal of the growth phase and rice varieties using sentinel-1a radar imagery in ciasem district, subang regency", IOP Conference Series: Earth and Environmental Science, 2022

Publication

<1 %

26

Zuhud Rozaki, Hermanto Siregar, Ikhsan Adi Pratama, Eni Istiyanti. "Food security, diversification, and inequality: Indonesia in the era of economic recovery and high price trends", Elsevier BV, 2023

Publication

<1 %

27

Moragoda Kankanamalage Jayani Chath Rajatheja, Rohana Chandrajith, Amita Bentota, Guttila Yugantha Jayasinghe. "A Comparative Assessment of Trace Element Accumulation in Native and Improved Rice (Oryza sativa L.) Varieties Grown Under Different Conditions of Fertilizer Application", Biological Trace Element Research, 2020

Publication

<1 %

28

N. K. Fageria. "Yield Physiology of Rice",
Journal of Plant Nutrition, 2007

Publication

<1 %

29

Sabrina J. Ruis, Charles Burr, Humberto
Blanco-Canqui, Brian Olson et al. "Corn
Residue Baling and Grazing Impacts on Corn
Yield under Irrigated Conservation Tillage
Systems", Agronomy Journal, 2021

Publication

<1 %

30

Divya Balakrishnan, Malathi Surapaneni,
Venkateswara Rao Yadavalli, Krishnam Raju
Addanki et al. "Detecting CSSLs and yield
QTLs with additive, epistatic and
QTL×environment interaction effects from
Oryza sativa × O. nivara IRGC81832 cross",
Scientific Reports, 2020

Publication

<1 %

31

Hanisa Sismaya Lestari, R. Selfi Nendris
Sulistiawan, Neng Siti Nurwasilah.
"IMPORTANCE PERFORMANCE ANALYSIS
KEPUASAN KONSUMEN PADA KUALITAS
PRODUK CABAI RAWIT DI PASAR
TRADISIONAL MUKA CIANJUR", AGRITA
(AGri), 2022

Publication

<1 %

32

Alakhyar Alakhyar, Fahrurrozi Fahrurrozi,
Widodo Widodo, Dia Novita Sari. "USE OF
GLIRICIDIA-ENRICHED LIQUID ORGANIC

<1 %

FERTILIZER FOR PRODUCTION OF CAISIM
(Brassica juncea L.)", Jurnal Agroqua: Media
Informasi Agronomi dan Budidaya Perairan,
2019

Publication

33

Mithra-Christin Hajati, Edwin Sutanudjaja, Nils Moosdorf. "Quantifying Regional Fresh Submarine Groundwater Discharge With the Lumped Modeling Approach CoCa-RFSGD", Water Resources Research, 2019

<1 %

Publication

34

D Roeswitawati, A Sulistyowati, E Ishartati. "Concentration Test of Liquid Bioslurry and Mushroom Baglog Waste as a Plant Medium on Growth and Results of Strawberries (Fragaria sp.)", IOP Conference Series: Earth and Environmental Science, 2021

<1 %

Publication

35

Irina Korotkova, Mykola Marenych, Volodymyr Hanhur, Oksana Laslo, Oksana Chetveryk, Viktor Liashenko. "Weed Control and Winter Wheat Crop Yield With the Application of Herbicides, Nitrogen Fertilizers, and Their Mixtures With Humic Growth Regulators", Acta Agrobotanica, 2021

<1 %

Publication

36

Muhammad Adnan, Koko Tampubolon, Fazal Ur Rehman, Muhammad Sulaman Saeed et

<1 %

al. "Influence of Foliar Application of Magnesium on Horticultural Crops: A Review", Agrinula : Jurnal Agroteknologi dan Perkebunan, 2021

Publication

37

Qasim Raza, Zulfiqar Ali, Ihsan Karim, Muhammad Ajmal, Muhammad Umer Khan. "Genetic Analysis of Triple Pistil Wheat Derived F2 Populations to Enhance Genetic Yield Potential", Research in Plant Biology, 1970

Publication

38

B Kurniasih, N Arini, D Alvioliana, R I Nisa, R A Wulandari. "Salt-induced growth promotion in rice varieties during nursery", IOP Conference Series: Earth and Environmental Science, 2021

Publication

39

Ranil Kavindra Asela Kularatne. "Sri Lanka's legal framework for marine pollution control: an evaluation of the Marine Pollution Prevention Act, No. 35 of 2008", Journal of the Indian Ocean Region, 2020

Publication

40

Shen, Yixin, Huixin Jiang, Guiyu Zhai, and Qingsheng Cai. "Effects of cutting height on shoot regrowth and forage yield of alfalfa (Medicago sativa L.) in a short-term cultivation system", Grassland Science, 2013.

Publication

<1 %

<1 %

<1 %

<1 %

41 Donald T. Krizek, Jerry C. Ritchie, Ali M. Sadeghi, Charles D. Foy, Errol G. Rhoden, John R. Davis, Mary J. Camp. "A Four-Year Study of Biomass Production of Eastern Gamagrass Grown on an Acid Compact Soil", Communications in Soil Science and Plant Analysis, 2011

<1 %

Publication

42 Erma Suryani, Rully Agus Hendrawan, Isnaini Muhandhis, Rarasmaya Indraswari. "A simulation model to improve the value of rice supply chain (A case study in East Java – Indonesia)", Journal of Simulation, 2020

<1 %

Publication

43 F.C. Oad ., Pompe Sta. Cruz ., N. Memon ., N.L. Oad ., Zia-Ul-Hassan .. "Rice Ratooning Management", Journal of Applied Sciences, 2001

<1 %

Publication

44 ELISANE GALBE DE CARVALHO COSTA, ALBERTO BAÊTA DOS SANTOS, FRANCISCO JOSÉ PFEILSTICKER ZIMMERMANN. "Crescimento da cultura principal e da soca de genótipos de arroz irrigado por inundação", Pesquisa Agropecuária Brasileira, 2000

<1 %

Publication

45

Tri Pujiana, Kordiyana K Rangga, Yuniar A Syarief, Abdul Mutolib. "Strategi Pengembangan Program Kawasan Rumah Pangan Lestari (KRPL) di Kabupaten Tulang Bawang Barat", Jurnal Ilmiah Membangun Desa dan Pertanian, 2020

Publication

<1 %

46

Publication

<1 %

47

Md. Abu Sayem Jiku, Md. Alimuzzaman, Ashutus Singha, Md. Arifur Rahaman et al. "Response and productivity of Garlic (*Allium Sativum* L.) by different levels of potassium fertilizer in farm soils", Bulletin of the National Research Centre, 2020

Publication

<1 %

48

Muslimin, Warda Halil, Wanti Dewayani. "Farmer's response to high yielding rice varieties in South Sulawesi (case study of Takalar)", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

49

Elham Yusefi-Tanha, Sina Fallah, Mohammad Pessarakli. " Effects of seed priming on growth and antioxidant components of hairy vetch () seedlings under chilling stress ", Journal of Plant Nutrition, 2019

Publication

<1 %

50

Kuster, C.. "Long-term trends in subsistence fishing patterns and coral reef fisheries yield from a remote Fijian island", Fisheries Research, 200511

Publication

<1 %

51

Yaer Karenius Mandacan, Detia Tri Yunandar, Susanti Indriya Wati. "Penyuluhan Pembuatan dan Pemanfaatan Pupuk Organik Cair Dikelompok Tani Sougp Hatam 1 Kampung Lismaunggu Distrik Prafi, Kabupaten Manokwari Provinsi Papua Barat", Prosiding Seminar Nasional Pembangunan dan Pendidikan Vokasi Pertanian, 2020

Publication

<1 %

52

A Hairani, M Noor. "Water management on peatland for food crop and horticulture production: research review in Kalimantan", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

53

Chang Zheng, Yuechao Wang, Desheng Yang, Sen Xiao, Yating Sun, Jianliang Huang, Shaobing Peng, Fei Wang. "Biomass, Radiation Use Efficiency, and Nitrogen Utilization of Ratoon Rice Respond to Nitrogen Management in Central China", Frontiers in Plant Science, 2022

Publication

<1 %

54

Eyitayo A. Makinde, Olukemi T. Ayoola, Esther A. Makinde. "Intercropping Leafy Greens and Maize on Weed Infestation, Crop Development, and Yield", International Journal of Vegetable Science, 2009

Publication

<1 %

55

L Z Nasution, C Hanum, Delvian. "The roles of FMA in increasing growth and effectiveness of soybean roots on dry land supporting sustainable agriculture in Deli Serdang District", IOP Conference Series: Earth and Environmental Science, 2020

Publication

<1 %

56

Mao Ye, Yuan-Yuan Song, Scott R. Baerson, Jun Long, Jie Wang, Zhiqiang Pan, Wen-Xiong Lin, Ren-Sen Zeng. "Ratoon rice generated from primed parent plants exhibit enhanced herbivore resistance", Plant, Cell & Environment, 2017

Publication

<1 %

57

U Susanto, C Gunarsih, W R Rohaeni. "Interaction of Genetic and Zn Fertilizer Application on Rice Yield and Grain Zinc Content", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

58

A Akter, MJ Hasan, MA Latif, MU Kulsum, PL Biswas, MH Rahman, RR Majumder, LF Lipi,

<1 %

MR Quddus, F Akter, A Ara. "Genetic Variability, Heritability, Correlation and Path Coefficient Studies for Yield and Yield Components of Some Promising Rice Hybrids", Bangladesh Rice Journal, 2020

Publication

59

A. K. Srivastava, Shyam Singh. "Citrus Decline: Soil Fertility and Plant Nutrition", Journal of Plant Nutrition, 2009

Publication

<1 %

60

Pawel Wójcik, Mariusz Lewandowski. "Effect of Calcium and Boron Sprays on Yield and Quality of "Elsanta" Strawberry", Journal of Plant Nutrition, 2003

Publication

<1 %

61

Prakash Jai, Singh Kamal. "Control of root-knot nematode by using composted sawdust in tomato root", African Journal of Biotechnology, 2014

Publication

<1 %

62

A. Ogunbayo S., K. Ojo D., A. Sanni K., G. Akinwale M. et al. "Genetic variation and heritability of yield and related traits in promising rice genotypes (Oryza sativa L.)", Journal of Plant Breeding and Crop Science, 2014

Publication

<1 %

63

A. Satyanarayana, T. M. Thiyagarajan, Norman Uphoff. "Opportunities for water saving with higher yield from the system of rice intensification", Irrigation Science, 2006

Publication

<1 %

64

AM Kartina, Susiyanti, W A Rusadi. "Characterization of Sugar Plant Seeds (*Arenga pinnata* Merr.) from Several Locations in Banten Province and Their Growth Response to the Provision of Liquid Organic Fertilizers", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

65

Arnita Chaorlina, Maryanti Setyaningsih, Hilman Faruq. "The Utilization of Tofu Waste Water as An Addition of Nutrition in Hydroponic Media to Lettuce Growth (*Lactuca sativa* L.)", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

66

Aynur Demir, Fulya Aydin-Kandemir. "Evaluation of Climate Change Impacts on the Geographic Distribution of *Fritillaria imperialis* L. (Liliaceae) (Turkey)", Acta Societatis Botanicorum Poloniae, 2022

Publication

<1 %

67

Daniela Ribeiro da Costa, Suzany Aguiar Leite, Vanderlúcia Fonseca de Paula, Jeferson

<1 %

Chagas do Nascimento et al. "Botanical insecticides and extracts and their lethal and sublethal effects on *Anastrepha obliqua* (Macquart) (Diptera: Tephritidae) and *Diachasmimorpha longicaudata* (Ashmead) (Hymenoptera: Braconidae)", Research Square Platform LLC, 2023

Publication

68

Dermiyati, E Aprilia, K Hendarto, S B Yuwono, Y C Ginting. "Effectiveness of the combination of organonitrofos and inorganic fertilizers on soil chemical properties and the yields of cucumber (*Cucumis sativus* L.) in Ultisols", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

69

Jason A. Bond, Patrick K. Bollich. "Ratoon Rice Response to Nitrogen Fertilizer", cm, 2006

Publication

<1 %

70

Mieczysław Czekański, Wojciech Antkowiak, Krzysztof Borkowski. "Adventitious Roots in Decaying Trunk of English Oak *Quercus robur* L.", *Wiadomości Botaniczne*, 2020

Publication

<1 %

71

P N Sari, M Auliya, U Farihah, N E A Nasution. "The effect of applying fertilizer of moringa leaf (*Moringa oliefera*) extract and rice washing water to the growth of pakcoy plant

<1 %

72

Sagrilo, Edvaldo, Pedro Soares Vidigal-Filho, Manoel Genildo Pequeno, Carlos Alberto Scapim, Maria Celeste Gonçalves-Vidigal, Rog rio Rui Maia, and Marcus Vin cius Kvitschal. "Efeito da  poca de colheita no crescimento vegetativo, na produtividade e na qualidade de ra zes de tr s cultivares de mandioca", Bragantia, 2002.

Publication

<1 %

73

Yunida Berliana, Juli Mutiara Sihombing, Khairani Khairani, Erfan Wahyudi. "The Influence of Cutting Age and Liquid Organic Fertilizer Dosage on The Yield of King Grass (Pennisetum purpupoides Schumach) as Source of Livestock", Agrinula : Jurnal Agroteknologi dan Perkebunan, 2021

Publication

<1 %

74

Hoi-Shun Lui, , M. Persson, and N. V. Z. Shuley. "Joint Time-Frequency Analysis of Transient Electromagnetic Scattering from a Subsurface Target", IEEE Antennas and Propagation Magazine, 2012.

Publication

<1 %

75

Lia Resti Utami, Yulian Yulian, Bambang Sulisty. "PERTUMBUHAN VEGETATIF BIBIT

<1 %

JERUK GERGA PASCA OKULASI PADA
KONSENTRASI PUPUK ORGANIK CAIR YANG
BERBEDA", Jurnal Ilmu-Ilmu Pertanian
Indonesia, 2019

Publication

76

Anan Polthane, Nisit Kumla, Bubpha Simma. "Effect of Pistia stratiotes, cattle manure and wood vinegar (pyroligneous acid) application on growth and yield of organic rainfed rice", Paddy and Water Environment, 2014

Publication

<1 %

77

C. Brondani, P. Rangel, R. Brondani, M. Ferreira. "QTL mapping and introgression of yield-related traits from *Oryza glumaepatula* to cultivated rice (*Oryza sativa*) using microsatellite markers", Theoretical and Applied Genetics, 2002

Publication

<1 %

78

E B M Adi, D Priadi, P Deswina, N W S Agustini. "The Growth of Pak Choy (*Brassica rapa* L.) on the microalgae (*Spirulina platensis*) biomass-based nutrient solution", IOP Conference Series: Earth and Environmental Science, 2023

Publication

<1 %

79

Muhammad Jamal Khan, Muhammad Hafeez Khan, Riaz Ahmad Khattak, Muhammad Tariq Jan. "Response of Maize to Different Levels of

<1 %

Sulfur", Communications in Soil Science and Plant Analysis, 2006

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off