

THE EFFECT OF PRUNING AND CHICKEN MANURE ON VEGETATIVE GROWTH OF HONEY DELI (*Syzygium aqueum* Burn F.) IN 9 MONTHS AGE

by Alridiwersah Alridiwersah

Submission date: 12-Oct-2023 07:48AM (UTC+0700)

Submission ID: 2192993845

File name: Growth_Of_Honey_Deli_Syzygium_aqueum_Burn_F_In_9_Months_Age.pdf (386.11K)

Word count: 6606

Character count: 32751

THE EFFECT OF PRUNING AND CHICKEN MANURE ON VEGETATIVE GROWTH OF HONEY DELI (*Syzygium aqueum* Burn F.) IN 9 MONTHS AGE

Alridiwersah¹, Rada Mulia Lubis², Aisar Novita¹

¹⁾ Lecturer in Department of Agrotechnology, Faculty of Agriculture, University of Muhammadiyah Sumatera Utara, Jl. Muchtar Basri No.3 Medan, 20238, Indonesia

²⁾ College Student in Department of Agrotechnology, Faculty of Agriculture, University of Muhammadiyah Sumatera Utara, Jl. Muchtar Basri No.3 Medan, 20238, Indonesia

Correspondence author: alridiwersah@umsu.ac.id

ABSTRACT

The study was conducted in April 2018 until June 2018, on the experimental field of the Faculty of Agriculture, Muhammadiyah University of North Sumatra, Jl. Tuar No.65 Medan Amplas District, Medan. Place height $\pm$ 27 masl. This study aims to determine the effect of pruning and administration of chicken manure on the vegetative growth of deli honey (*Syzygium aqueum* Burn F.) at the age of 9 months. This study uses Factorial Randomized Block Design (RBD) with 2 treatment factors, namely Pruning with 3 levels, namely P: Without Pruning, P: 75 cm, P: 85 cm and Chicken Manure with 4 levels, namely A: control, A: 100 gram / polybag, A: 200 gram / poly bag, A: 300 gram / polybag. The parameters measured were plant height, stem diameter, leaf size, number of branches, leaf area, age of flowering, leaf chlorophyll. The results showed that pruning treatment had a significant effect on the parameters of the height of honey medicinal plants and the administration of chicken manure significantly affected stem diameter, number of leaves, number of branches, leaf area and age, and the interaction between pruning and chicken manure had no significant effect on all parameters observation.

Keywords: guava honey, pruning, chicken manure.

A. INTRODUCTION

Guava (*Syzygium aqueum* Burn F. Alston) originates from the Indo-China and Indonesia regions spread to Malaysia and the islands in the Pacific. So far, it is still concentrated as a garden plant for family consumption. Guava water is not just sweet refreshing, but has diversity in appearance. Water guava is categorized as one of the potential fruit types that has not been widely cultivated for commercial purposes. Its perishable nature is an important problem that needs to be solved. The fruit can be said to be physically damaged skin so little on the fruit will accelerate rot on the green deli water buah. Jambu crops introduced into varieties that have been released in 2012. Based on the research that guava green deli has a water content of 81.596%, levels sugar 12.4°brix, vitamin C 210.463 mg/100g and has a sweet taste like honey. This guava grows well at altitudes of 0-500 meters above sea level (Chairani et al., 2015).

Problems that are often faced in deli honey guava cultivation include plants that are late to bear fruit even though their age is supposed to produce. This is due to the lack of nutrients needed by deli honey guava plants in their growth so that production is not optimal. Besides that, the size and taste of fruit is often a problem, but among farmers there are still many who do not know how to get deli honey fruit with optimal size and taste. So often the production of guava honey deli with a small size of fruit with a tasteless taste.

According to Balitbu (2015), the criteria for guava which are suitable for harvesting have

characteristics such as, having a fruit weight of 200-300 gr / seed and having a sweet range of 12-14 brix to obtain these criteria can be done by pruning and giving fertilizers that contains magnesium.

Fitria (2016), explained that pruning activities are activities to dispose of unproductive parts of plants so that new shoots can emerge that will produce fruit. The more crop branches that are pruned, the more productive shoots that emerge and produce fruit.

Pruning on guava honey is an attempt to improve environmental conditions such as temperature, humidity, light, wind circulation so that photosynthetic activity takes place normally. Pruning can improve plant health, flowering is aroused and production increases. There are several backgrounds that underlie why plants should be trimmed, ie plants tend to grow steadily, both grow upward and grow sideways. The benefits of top shoot pruning will reduce competition from photosynthesis between leaves and fruit and reduce the incidence of disease, as well as pruning on tomatoes, which increases fruit size. If excessive growth leaf shoots is trimmed, the circulation of air around the canopy improves, this condition will reduce the humidity of the microclimate around the plant and so will reduce the incidence of disease in addition to increasing the yield of flowers can also improve the quality of flowers and appearance or plant figures for the better but it can hold declining flower production (Saladin, 2002).

Chicken manure is a fertilizer that comes from a mixture of chicken manure. Organic material plays an important role in the formation of good and stable soil structures so that the infiltration and ability to store water. According to the provision of manure, it significantly reduces the amount of surface flow because manure improves the physical properties of the soil, especially the structure so that permeability increases.

The way to deal with deli honey guava which is late in the fruit can be done by providing fertilizer to meet the nutrients needed by deli guava in its growth such as providing chicken manure containing macro nutrients such as N, P, and K. In addition to providing fertilizer, pruning can also speed up the honey guava plant to bear fruit. This is because the results of plant photosynthesis are not only channeled for vegetative growth of plants but also for the generative growth of plants such as flowers and fruit, thereby accelerating the production of deli guava honey. Besides that, pruning efforts can also affect the size of the fruit produced by pruning the resulting fruit will be even greater. Meanwhile, to get guava honey fruit with a sweet taste can be done by providing fertilizers containing magnesium.

Based on the above problems, this study was conducted to determine the effectiveness of giving chicken manure and pruning to the production of deli guava honey. So that later this research can provide benefits for those who do guava honey cultivation.

B. MATERIALS AND METHODS

Place and time

This research was conducted in April 2018 until June 2018, on the experimental field of the Faculty of Agriculture, Muhammadiyah University

of North Sumatra II. Tuar No. 65 Medan Amplas District, Medan. Place height ± 27 masl.

Materials and tools

The materials used in this study were 9 months old green deli honey guava seeds, topsoil soil, chicken manure, water, insecticides, fungicides, polybags measuring 25 cm x 35 cm, plant signposts.

The tools used are hoes, bolts, meters, raffia ropes, machetes, knives, buckets, scissors, calculators, barrels/buckets, wood, cameras and stationery.

Research methods

This study used factorial randomized block design (RBD) with two factors studied, namely: Pruning Factor (P) with 3 levels (P_0 : Without pruning, P_1 : 75 cm, P_2 : 85 cm) and factors of giving solid chicken manure (A) with 4 levels (A_0 : control, A_1 : 100 gram/polybag, A_2 : 200 gram/polybag, A_3 : 300 gram/polybag).

The parameters measured were plant height, stem diameter, number of leaves, number of branches, leaf area, age and flowering leaf chlorophyll.

C. RESULTS AND DISCUSSION

Plant height

Based on the analysis of variance (ANOVA) with Randomized Block Design (RAK) showed that pruning had a significant effect on plant height parameters while giving chicken manure and the interaction of the two factors has no significant effect on plant parameters 2, 4, 6, and 8 WAP. The average plant height can be seen in tables 1 - 3.

Table 1. Average of Honey Guava Plant Height 2, 4, 6, and 8 WAP in Pruning Treatment.

Pruning	Age (WAP)			
	2	4	6	8
.....cm.....				
P_0 = Without pruning	120.92	125.17	130.29	135.29 a
P_1 = 75 cm	104.04	108.17	113.29	118.29 b
P_2 = 85 cm	103.42	108.67	112.88	118.13 bc

Note: Numbers followed by letters that are not the same in the same column are significantly different according to the 5% DMRT Test.

Table 2. Average height of guava plants aged 2, 4, 6 and 8 WAP in chicken manure fertilizer treatment.

Manure Chicken	Age (WAP)			
	2	4	6	8
.....cm.....				
A_0 = control	111.06	115.89	120.39	124.94
A_1 = 100 gram/polybag	106.94	110.28	116.61	121.61
A_2 = 200 gram/polybag	112.22	116.33	120.39	125.56
A_3 = 300 gram/polybag	107.61	113.50	117.89	123.50

Table 3. Average Interaction of Treatment for Pruning and Manure of Chicken at Honeycomb Plant Height 2, 4, 6, and 8 WAP.

Interaction	Age (WAP)			
	2	4	6	8
	cm.....			
P ₀ A ₀	130.17	135.17	140.17	144.83
P ₀ A ₁	114.50	115.17	124.00	128.83
P ₀ A ₂	122.00	123.33	125.17	130.33
P ₀ A ₃	117.00	127.00	131.83	137.17
P ₁ A ₀	102.00	104.50	109.50	113.83
P ₁ A ₁	103.50	108.00	113.17	118.17
P ₁ A ₂	112.33	118.00	123.33	128.33
P ₁ A ₃	98.33	102.17	107.17	112.83
P ₂ A ₀	101.00	108.00	111.50	116.17
P ₂ A ₁	102.83	107.67	112.67	117.83
P ₂ A ₂	102.33	107.67	112.67	118.00
P ₂ A ₃	107.50	111.33	114.67	120.50

Based on Table 1, it can be seen that the honey guava plant height with the highest average is found in the treatment P₀ (control) is 135.29 cm which is significantly different from the treatment P₁ (75 cm) which is 118.29 cm and P₂ (85 cm) which is 118.13 cm.

Relationship between honey guava plant height and treatment pruning can be seen in Figure 1.




Figure 1. Graph of Plant Height with Pruning Treatment.

Based on Figure 1 it can be seen that pruning with optimum length is 85 cm with the highest average of 135.29 cm which shows a negative linear relationship with the regression equation $\hat{y} = 188.28 + (-0.8583)x$ with $r = 0.7573$. Based on the equation, it can be seen that plant height will decrease along with the increase in length of pruning.

In the parameters of plant height it is known that shoot trimming can reduce the growth of plant height when compared with plants without pruning treatment. Pruning will result in reduced plant height due to organ removal. The removal of apical buds aims to stop the growth of shoots and help accelerate the growth of lateral shoots which give rise to new branches in the axillary main stem.

Pane, et al. (2013), stated that shoot trimming can suppress plant height. Plants that are not pruned have higher plant height while pruned plants have lower plant height. Plants that are not pruned will continue to grow because the hormone auxin in shoots is very high. Although the plants that were given pruning treatment had lower plant height, the plants had a greater number of branching when compared to plants that were not given shoot trimming treatment.

Trimming shoots can suppress the growth of apical shoots or shoots and maximize the growth of lateral shoots, so that the formation of new branches will be balanced and then have an impact on increasing plant productivity. Trimming shoots can be done by cutting the top of plants, branches

and leaves. According to Esrita (2012), that shoot trimming is proven to increase the number of productive branches on plants.

Stem Diameter

Based on the analysis of variance (ANOVA) with Randomized Block Design (RAK) showed

that the administration of chicken manure had a significant effect on stem diameter parameters while pruning and interaction of the two factors had no significant effect on plant parameters 2, 4, 6, and 8 WAP. The average number of branches can be seen in tables 4 – 6.

Table 4. Average Diameter of Honey Guava Age 2, 4, 6, and 8 WAP in Pruning Treatment.

Pruning	Age (WAP)			
	2	4	6	8
	cm.....			
P ₀ = Without pruning	2.35	2.45	2.55	2.69
P ₁ = 75 cm	2.29	2.40	2.51	2.65
P ₂ = 85 cm	2.33	2.42	2.55	2.68

Table 5. Average Diameter of Honeycomb Age 2, 4, 6, and 8 WAP in Chicken Cage Fertilizer Treatment.

Manure Chicken	Age (WAP)			
	2	4	6	8
	cm.....			
A ₀ = control	2.22	2.33	2.44	2.53cd
A ₁ = 100 gram/polybag	2.28	2.39	2.51	2.61bc
A ₂ = 200 gram/polybag	2.41	2.49	2.59	2.69b
A ₃ = 300 gram/polybag	2.38	2.48	2.62	2.85a

Note: Numbers followed by letters that are not the same in the same column are significantly different according to the 5% DMRT Test.

Based on Table5, it can be seen that the number of leaves guava honey with the highest average is found in the treatment A₃ (300 gram/ polybag) which is 2.85 cm which is significantly different from the

treatment A₂ (200 gram/ polybag) which is 2.69 cm, A₁ (100 gram/ polybag), which is 2.61 cm and A₀ (control) which is 2.53 cm

Table 6. Average Interaction of Pruning and Chicken Manure Treatment on 2, 4, 6, and 8 WAP Honeycomb Stem Diameter.

Interaction	Age (WAP)			
	2	4	6	8
	cm.....			
P ₀ A ₀	2.23	2.33	2.42	2.52
P ₀ A ₁	2.30	2.42	2.55	2.65
P ₀ A ₂	2.43	2.48	2.60	2.67
P ₀ A ₃	2.45	2.55	2.65	2.92
P ₁ A ₀	2.23	2.37	2.48	2.57
P ₁ A ₁	2.20	2.32	2.43	2.55
P ₁ A ₂	2.45	2.55	2.65	2.75
P ₁ A ₃	2.27	2.37	2.48	2.73
P ₂ A ₀	2.20	2.28	2.42	2.50
P ₂ A ₁	2.35	2.43	2.53	2.63
P ₂ A ₂	2.35	2.43	2.53	2.67
P ₂ A ₃	2.43	2.52	2.73	2.90

Relationship between the honey guava plant height and the administration of chicken manure can be seen in Figure 2.




Figure 2. Graph of Stem Diameter with Chicken Manure Fertilizer Treatment.

Based on Figure 2, it can be seen that the administration of chicken manure with the optimum dose is 300 grams/polybag with the highest average 2.85 cm which shows a positive linear relationship with the regression equation $\hat{y} = 2.5133 + 0.0011x$ with $r = 0.9724$. Based on this equation, it can be seen that the stem diameter will increase along with the increase in chicken manure dosage. Increasing vegetative growth in stem diameter parameters is strongly influenced by the role of nutrients such as N, P and K. Lingga and Marsono (2003) explain that the role of nitrogen in plants is to stimulate overall growth, especially branches, stems and leaves. Nitrogen functions as a form of chlorophyll, protein and fat. Nitrogen is also a constituent of enzymes contained in cells, thus affecting the growth of carbohydrates which play a role in plant growth.

Sosrosoedirdjo (2004) added that carbohydrates are a material that is very necessary in cell division, cell extension, cell enlargement and tissue formation for the development of stems, leaves and roots. Phosphorus functions to overcome the negative effects of nitrogen, improve root

development and improve the quality of results. Then K functions in regulating the balance of nitrogen and phosphorus fertilizers (Jumin, 2000).

Good growth is indicated by the ability of plants to photosynthesize higher and produce more photosynthesis (photosynthate). Photosynthate which is more translocated through phloem and can be used to stimulate secondary growth, namely expansion of stem cells and indicated by wider stem diameter. According to Gardner, et al. (1991) growth and development of plants and their organs will depend on the availability of meristems, hormones and photosynthesis (carbohydrates) and the supporting environment.

Number of leaves

Based on the analysis of variance (ANOVA) with Randomized Block Design (RAK) showed that administration of chicken manure had a significant effect on the parameters of the amount of dung while tripping and the interaction of the two factors has no significant effect on the parameters of number 2, 4, 6, and 8 WAP leaves. The average number of leaves can be seen in Tables 7-9.

Table 7. Average Amount of Honey Leaf 2, 4, 6, and 8 WAP in Pruning Treatment.

Pruning	Age (WAP)			
	2	4	6	8
	strands.....			
P ₀ = Without pruning	5.63	10.83	20.83	30.88
P ₁ = 75 cm	6.08	11.21	21.21	31.75
P ₂ = 85 cm	5.96	10.83	20.83	30.96

Table 8. Average Amount of Honey Leaf 2, 4, 6, and 8 WAP in Chicken Cage Fertilizer Treatment.

Manure Chicken	Age (WAP)			
	2	4	6	8
	 strands.....			
A ₀ = control	5.33	10.28	20.33	28.94 cd
A ₁ = 100 gram/polybag	5.44	10.50	20.50	30.44 bc
A ₂ = 200 gram/polybag	5.78	10.72	20.72	31.61b
A ₃ = 300 gram/polybag	7.00	12.33	22.28	33.78 a

Note: Numbers followed by letters that are not the same in the same column are significantly different according to the 5% DMRT Test.

Based on Table 8, it can be seen that thenumber of leavesguava honeywith the highest average is found in the treatment A₃(300gram/ polybag) which is 33.78 strands which is

significantly different from the treatment A₂ (200gram/polybag), which is 31.61 strands, A₁ (100gram/ polybag), which is 30.44 strands and A₀ (control), which is 28.94 strands.

Table 9. Average Interaction of Pruning and Chicken Manure Treatment in Amount of 2, 4, 6, and 8 WAP Guava Leaf Leaves.

Interaction	Age (WAP)			
	2	4	6	8
	 strands.....			
P ₀ A ₀	5.33	10.33	20,33	27.83
P ₀ A ₁	5.33	10.33	20.50	30.83
P ₀ A ₂	5.50	10.50	22,17	31,17
P ₀ A ₃	6.33	12,17	20.50	33.67
P ₁ A ₀	5.33	10.33	20.50	29.50
P ₁ A ₁	5.50	10.50	21.00	30,33
P ₁ A ₂	6.00	11.00	22.83	32.67
P ₁ A ₃	7.50	13.00	20,17	34.50
P ₂ A ₀	5.33	10.17	20.67	29.50
P ₂ A ₁	5.50	10.67	20.67	30,17
P ₂ A ₂	5.83	10.67	21.83	31.00
P ₂ A ₃	7.17	11.83	20,33	33.17

The relationship between the heigh tofguava honey and the administration of chicken manure can be seen in Figure 3.





Figure 3.Graph of Number of Leaves with Chicken Cage Fertilizer Treatment.



Based on Figure 3, it can be seen that the administration of chicken manure with optimum dosage is 300 grams/polybag with the highest average of 33.78 strands which shows a positive linear relationship with the regression equation $\hat{y} = 28.844 + 0.0157 x$ with $r = 0.984$. Based on these equations it can be seen that the number of leaves will increase with increasing doses of chicken manure.

Provision of chicken manure significantly affected the number of leaves of guava honey. This is because manure contains complete macro and micro nutrient. Nutrient content of N (2.71%) which is high in chicken manure stimulates plant growth in general. Nitrogen plays a role in the formation of chlorophyll, amino acids, fats and enzymes. While nutrient P (6.31%) plays a role in root growth and development. Element K (2.01%) helps the formation of proteins and minerals and increases plant resistance to disease (Purwa, 2009).

The presence of nitrogen elements will increase the growth of vegetative parts such as

leaves. This is in accordance with the opinion of Lingga and Marsono (2003), that the main role of nitrogen for plants is to stimulate overall growth, especially the stems, branches and leaves. High nitrogen content in chicken manure stimulates the growth rate of plant leaves. Sutedjo (2002) added that nitrogen is the main nutrient needed for the growth of vegetative parts of leaves, stems and roots, but if given excessively it can inhibit flowering and fertilization in plants.

Number of branches

Based on the analysis of variance (ANOVA) with Randomized Block Design (RAK) showed that administration of chicken manure significantly affected the parameters of the number of branches while pruning and the interaction of the two factors has no significant effect on the parameters of number 2, 4, 6, and 8 WAP branches. The average number of branches can be seen in tables 10-12.

Table 10. Average Number of Branches of Honey Guava Age 2, 4, 6, and 8 WAP in Pruning Treatment.

Pruning	Age (WAP)			
	2	4	6	8
	branches.....			
P ₀ = Without pruning	4.50	4.50	4.50	4.92
P ₁ = 75 cm	4.54	4.46	4.46	5.08
P ₂ = 85 cm	4.50	4.50	4.54	5.33

Table 11. Average Interaction of Chicken Cage Pruning and Fertilizer Treatment in Number of 2, 4, 6, and 8 WAP Guava Branches.

Interaction	Age (WAP)			
	2	4	6	8
	branches.....			
P ₀ A ₀	4.17	4.17	4.17	4.17
P ₀ A ₁	4.00	4.00	4.17	4.50
P ₀ A ₂	4.17	4.17	4.00	4.67
P ₀ A ₃	5.67	5.67	5.67	6.33
P ₁ A ₀	4.00	4.00	4.00	4.17
P ₁ A ₁	4.33	4.33	4.33	4.83
P ₁ A ₂	4.33	4.33	4.33	5.17
P ₁ A ₃	5.50	5.17	5.17	6.17
P ₂ A ₀	4.17	4.17	4.33	4.67
P ₂ A ₁	4.33	4.33	4.33	5.00
P ₂ A ₂	4.00	4.00	4.00	5.17
P ₂ A ₃	5.50	5.50	5.50	6.50

Based on Table 11, it can be seen that the number of branches guava honey with the highest average is found in the treatment A₃ (300 gram/polybag) which is 6.33 branches which is

significantly different from the treatment A₂ (200 gram/polybag) which is 5.00 branches, A₁ (100 gram/polybag) which is 4.78 branches and A₀ (control) which is 4.33 branches.

Table 12. Average Number of Branches of Honey Guava Age 2, 4, 6, and 8 WAP in Chicken Manure Fertilizer Treatment.

Manure Chicken	Age (WAP)			
	2	4	6	8
	branches.....			
A ₀ = control	4.11	4.11	4.17	4.33cd
A ₁ = 100 gram/polybag	4.22	4.22	4.28	4.78bc
A ₂ = 200 gram/polybag	4.17	4.17	4.11	5.00b
A ₃ = 300 gram/polybag	5.56	5.44	5.44	6.33a

Note: Numbers followed by letters that are not the same in the same column are significantly different according to the 5% DMRT Test.

Relationship between thenumber of branches guava honey by giving chicken manure can be seen in Figure 4.




Figure 4. Graph of Number of Branches with Chicken Manure Fertilizer Treatment.

Based on Figure 4, it can be seen that the administration of chicken manure with the optimum dose is 300 grams / polybag with the highest average 6.33 branches which show a positive linear relationship with the regression equation $\hat{y} = 4.1778 + 0.0062x$ with $r = 0.8711$. Based on these equations it can be seen that the number of branches will increase with increasing doses of chicken manure.

Fertilizer Chicken coops have an influence on the number of branches because chicken manure has a high N content, so that more branches are formed. Increased value of vegetative characters such as plant height and number of branches caused by the role of nitrogen elements. The main role of nitrogen for plants is to stimulate overall growth, especially the stems, branches, and leaves (Hardjowigeno, 2007). This is in accordance with the opinion of Gardner (1991), that in addition to environmental factors such as water availability, nutrient deficiencies in the soil can also affect

vegetative growth. Sutedjo and Kartasapoetra (1987), said that plants need nutrients that are suitable for their needs in the process of growth and development.

According to Suryanto (1999), the availability of sufficient nitrogen is needed to support plant growth and development. Elements of nitrogen absorbed by plants in sufficient amounts will stimulate meristematic tissue at the point where the stem grows more active. This can spur plant growth.

Leaf area

Based on the analysis of variance (ANOVA) with Randomized Block Design (RAK) showed that administration of chicken manure significantly affected leaf area parameters while pruning and the interaction of the two factors has no significant effect on leaf area parameters 2, 4, 6, and 8 WAP. The average leaf area can be seen in tables 13-15.

Table 13. Average Area of Honey Guava Leaf Age 2, 4, 6, and 8 WAP in Pruning Treatment.

Pruning	Age (WAP)			
	2	4	6	8
	cm ²			
P ₀ = Without pruning	44.33	77.56	124.00	168.44
P ₁ = 75 cm	46.65	77.19	127.19	173.68
P ₂ = 85 cm	47.01	70.09	124.13	173.19

Table 14. Average Area of Honey Guava Leaf Age 2, 4, 6, and 8 WAP in Chicken Cage Fertilizer Treatment.

Treatment Manure Chicken	Age (WAP)			
	2	4	6	8
	cm ²			
A ₀ = control	41.13	73.84	116.01	163.70bcd
A ₁ = 100 gram/polybag	41.94	74.32	116.62	165.53bc
A ₂ = 200 gram/polybag	44.61	74.22	124.88	172.21b
A ₃ = 300 gram/polybag	56.31	77.39	142.91	185.63a

Note: Numbers followed by letters that are not the same in the same column are significantly different according to the 5% DMRT Test.

Based on Table 14, it can be seen that the number of leaves guava honey with the highest average is found in the treatment A₃(300 gram/polybag) which is 185.63 cm² which is significantly different from the treatment A₂ (200gram/ polybag) which is 172.21 cm², A₁ (100gram/ polybag), is 165.53 cm² and A₀(control), which is 163.70 cm².

Table 15. Average Interaction of Treatment of Pruning and Chicken Manure in the Area of 2, 4, 6, and 8 WAP Guava Leaves.

Interaction	Age (WAP)			
	2	4	6	8
	cm ²			
P ₀ A ₀	41.19	73.48	115.50	161.74
P ₀ A ₁	41.19	73.19	117.07	164.40
P ₀ A ₂	44.05	77.18	133.22	170.43
P ₀ A ₃	50.89	86.38	130.20	177.18
P ₁ A ₀	41.31	75.83	117.13	164.99
P ₁ A ₁	41.26	73.65	114.84	165.05
P ₁ A ₂	48.79	83.51	127.22	179.93
P ₁ A ₃	55.24	75.79	149.55	184.76
P ₂ A ₀	40.89	72.22	115.40	164.38
P ₂ A ₁	43.38	76.13	117.93	167.15
P ₂ A ₂	40.99	61.99	114.19	166.28
P ₂ A ₃	62.80	70.00	148.98	194.95

Relationship between leaf area guava honey by providing chicken manure can be seen in Figure 5.




Figure 5. Wide Leaf Graph with Chicken Cage Fertilizer Treatment.

Based on Figure 5, it can be seen that the administration of chicken manure with the optimum dose is 300 grams / polybag with the highest average 185.63 cm² which shows a linear positive relationship with the regression equation $\hat{y} = 160.9 + 0.0725x$ with $r = 0.8862$. Based on these equations, it can be seen that leaf area will increase with increasing doses of chicken manure.

Chicken manure given to guava honey has a significant influence on leaf area parameters. This is because nutrients in chicken manure can be absorbed by plants for the process of division, cell enlargement and differentiation which causes additional volume in the leaves of plants. Nutrients that play an important role in this case are nutrients N, P and K. This is in accordance with the statement Damanik et al (2009) which states that nutrient N is very instrumental in the formation of plant cells, tissues and plant organs. N element is very necessary when plants enter the vegetative growth phase. P nutrients play an important role in absorbing plants against nutrients in the soil. K

nutrients play a major role in photosynthesis and carbohydrate translocation and also regulate the distribution of water in plants. Lack of K elements will cause the leaves to fall.

Erawan (2013) added that the increase in leaf area of plants was due to the availability of nitrogen elements that can be absorbed by plants, so that plant growth such as leaf area also increased. The provision of chicken manure has the highest yield which shows that chicken manure can provide better nutrients than other manure.

Age begins to flower

Based on the analysis of variance (ANOVA) with Randomized Block Design (RAK) showed that administration of chicken manure had a significant effect on age parameters starting to flower while pruning and the interaction of the two factors has no significant effect on age parameters starting to flower. The average age of flowering can be seen in table 16

Table 16. Average age begins to flower guava in chicken cage pruning and fertilizer treatment.

Pruning	Chicken Manure				Average
	A ₀	A ₁	A ₂	A ₃	
	WAP				
P ₀ = Without pruning	8.33	8.00	8.00	6.50	7.71
P ₁ = 75 cm	8.50	8.33	8.00	7.00	7.96
P ₂ = 85 cm	8.33	8.00	7.50	7.17	7.75
Average	8.39cd	8.11bc	7.83b	6.89a	

Description: Numbers followed by letters that are not the same on the same line are significantly different according to the 5% DMRT Test.

Based on Table 16, it can be known that guava honey starts flowering with the average tar contained in the treatment A₃ (300 gram/polybag) which is 6.89 WAP which is significantly

different from the treatment A₂ (200 gram/polybag) which is 7.83 WAP, A₁ (100 gram/polybag) which is 8.11 WAP and A₀ (control) which is 8.39 WAP.

The relationship between age begins to flower guava honey by providing chicken manure can be seen in Figure 6.




Figure 6. Age Chart Begins to Bloom with the Treatment of Chicken Cage Fertilizer.

Based on Figure 6, it can be seen that the administration of chicken manure with the optimum dose is 300 grams/polybag with the highest average 6.89 which shows a positive linear relationship with the regression equation $\hat{y} = 8.5222 + (-0.0048)x$ with $r = 0.8954$. Based on these equations, it can be seen that the age of flowering will be faster with increasing doses of chicken manure.

Provision of chicken manure has an effect on the age of flowering of guava honey. This is because the speed of flowering is influenced by the availability of nutrients in the soil. Soil that is given chicken manure can add nutrients. Gardner et al. (1991), states that there are two factors that affect the speed of flowering in plants, first an external factor (environment), namely sunlight that plays an important role in the progress of photosynthesis, when sunlight is appropriate it will affect the speed of flowering of a plant and the availability of nutrients land that is related to the availability of energy supply and building materials for the

formation and development of flowers. The second is internal factors (genetic), if the minimum age is met, the plants will flower.

According to Salisbury and Ross (1995) the age of flowering or when the first flower appears from a variety planted at the same time and environment, the possibility of age of flowering in plants is almost the same. Temperature is one of the environmental factors that influences flower formation. Temperature directly affects photosynthesis, respiration, transpiration, water absorption and nutrients.

Leaf chlorophyll

Based on the analysis of variance (ANOVA) with randomized block design (RBD) showed that pruning and administration of chicken manure and the interaction of the two factors had no significant effect on leaf chlorophyll parameters. The average chlorophyll of leaves can be seen in table 17.

Table 17. Average of Guava Leaf Chlorophyll in Chicken Cage Pruning and Fertilizer Treatment

Pruning	Chicken Manure				Average
	A ₀	A ₁	A ₂	A ₃	
	mg/g				
P ₀ = Without pruning	42.37	43.08	40.48	33.60	39.88
P ₁ = 75 cm	41.00	43.55	38.77	46.15	42.37
P ₂ = 85 cm	45.08	42.33	43.83	43.47	43.68
Average	42.82	42.99	41.03	41.07	

Based on table 17, it can be seen that chlorophyll leaves with the highest average of pruning treatment were found in treatment P₂ (85 cm), which was 43.68 mg/g and the lowest in

treatment P₀ (control) was 39.88 mg/g. While chlorophyll leaves with the highest average treatment of chicken manure are found in A₁ (100 grams/polybag), which is 41.07 mg/g and the

lowest is in treatment A₂ (200 grams/polybag), which is 41.03 mg/g. Of the two treatments, there was no significant effect and interaction on leaf chlorophyll.

D. CONCLUSIONS AND SUGGESTIONS

Conclusion

Based on the results of analysis of experimental data in the field, it can be concluded as follows:

1. The pruning treatment has a significant effect on the parameters of the honey guava plant height.
2. Provision of chicken manure significantly affected stem diameter, number of leaves, number of branches, leaf area and age began to flower.
3. The interaction between pruning and chicken cage fertilizer did not significantly affect all observational parameters.

Suggestion

Based on the results of the study it is recommended that further research be conducted by using higher doses of chicken manure and higher pruning as well as on different locations to obtain maximum results on the cashew crop commodity.

REFERENCES

- Barus, A. and Syukri. 2008. Agrotechnology of Fruits. USU Press, Medan. 147-150 p.
- Cahyono, B. 2010. Success in Cultivating Water Guava in Farms and Estates. Andy: Yogyakarta.
- Central Bureau of Statistics [BPS]. 2013. Annual Production of Fruits and Vegetables in Indonesia. [Internet] [downloaded 2014 Dec 27].
- Chairani, H., Revandy IMD and Victor, HT 2015. Vegetative and Generative Growth of Guava (*Syzygium samarangense* Blume Merr and Perry) Green Deli Varieties with ZPT Treatment and Planting Media. Online Journal of Agroecotechnology. ISSN No. 2332-6597 Vol. 3.3, No. 2: 740-747, March 2015.
- Damanik, MMB, Hasibuan, BE, Izzu, Sarifuddin and Hanum, H. 2009. Soil Fertility and Fertilization. USU press. Field.
- Erawan, D., WO Yani, and A. Bahrin. 2013. Growth and yield of mustard (*Brassicajunceal.*) plants at various doses of urea fertilizer. J. Agroteknos. 3 (1): 19-25.
- Esrita, D. 2012. Effect of Apical Pruning on Soybean Growth and Yield (*Glycine max L.*). J. Bioplantae. 1 (2): 125-133.
- Fitria, L. 2016. Study of Growth, Production and Quality of Guava (*Psidium guajava L.*) Var. Crystals in the Origin of Seeds and Different Pruning. Thesis of Bogor Agricultural Institution Graduate Program.
- Gardner, FP, RB Pearce and RI Mitchell. 1991. Cultivation Physiology, UI Press Jakarta.
- Hardjowigeno, S. 2007. Soil Science. Jakarta: Akademika Pressindo.
- Hartawan, R., 2008. Variability of Water Guava Seed Growth. Origin of Superior and Wild Seeds. Journal of Media Academic Vol. 2 No.1 pp. 34-43.
- Herdiantoro, D. 2013. Factorial Design (Completely Randomized Design & Randomized Block Design). Soil Biology Laboratory, Soil Science Department, Faculty of Agriculture. Padjadjaran University: Bandung.
- Iriani, NM, Sofiyanti N. and Fitmawati. 2014. Analysis of the relationship of guava water (*Syzygium aqueum* (Burm.f.). Alston) in Pekanbaru and Kampar District based on morphological characters. JOM FMIPA. 1 (2): 1-7.
- Islami, T. 1997. Relations of Land, Water and Plants. IKIP Semara Press. Semarang. Items 234-244.
- Jumin. 2000. Basics of Agronomy. Rajawali. Bandung.
- Kartasapoetra AG, Sutedjo MM. 1987. Fertilizer and Fertilization Method. Jakarta: Rineka Cipta.
- Lakitan, B. 2005. Physiology of Plant Growth and Development. PT Raja Grafindo Persada. Jakarta. hlm 219-229.
- Lingga, P., and Marsono. 2003. Directions for Using Fertilizers. Self Help Spreader. Jakarta.
- Margianasari AF, Junaedi, Sugono J and Zen D. 2013. Practical Guides for Planting Fruits on Land and Pot. Jakarta (ID): Penebar Swadaya.
- Pane, SI, L. Mawarni and T. Irmansyah. 2013. Response of Soybean Growth to Pruning and Provision of OPEFB Compost on Shaded Land. Online Journal of Agroecotechnology. 1 (2): 393-401.
- Purwa, DR, 2009. Instructions for Fertilization. AgroMedia Pustaka, Jakarta.
- Saladin, M. 2002. Evaluation of Horticulture Character and the Effect of Branch Pruning on Production of Ten Tomatoes of Hope of IPB. Essay. Agriculture Cultivation Department. Faculty of Agriculture. Bogor Agricultural Institute. Bogor.
- Salisbury, FB and Ross CW 1995. Plant Physiology Volume 2. Bandung Press Institute of Technology. Bandung.
- Shinta, DA 2016. Characterization of Morphology and Anatomy of Water Guava Plants in the Faculty of Mathematics and Natural

- Sciences.Essay.Mekarsari Bogor
Agricultural Institute Bogor, West Java.
- Soekartawi, 1995. Analysis of Farming for Smallholder Development, University of Indonesia.
- Sosrosoedirdjo. 2004. Science of Fertilizing. Volume I. CV. Yasaguna. Jakarta.
- Sri, 2002. Effects of Water Stress on Plant Physiology. IPB Press. Bandung. Pp. 208-218.
- Sumenda, L. 2011. Analysis of the Content of Mango Leaf Chlorophyll (*Mangifera Indica* L.) at Different Leaf Development Levels. *Bioslogos*, pp. 1-10.
- Suryanto, A. 1999. Study of the Form and Dosage of Nitrogen Fertilizers in Broccoli Plants. *J. Habitat*. 10 (108): 1. - 9
- Sutedjo, MM, and AG Kartasaputra, 1990. *Fertilizer and Fertilization*. Rineka Cipta, Jakarta.
- Tropical Fruit Research Center. 2015. Prospects for Accurate Cultivation of Honey Guava Deli. Biology Education Study Program, Teacher Training and Education Faculty of Surakarta Muhammadiyah University.
- 14 Tsukaya H. 2005. Leaf Shape: Genetic Controls And Environmental Factors. *Int J Dev Biol*. 49: 547-555.1.
- UPT. (BPSB) Binjai Central Bureau of Statistics. 2012. Proposal for Registration of Honey Deli Water Guava Varieties (Origin of Binjai City). Department of Agriculture of North Sumatra Province. Field
- Verheij, EWM and RE Coronel, 1997. Southeast Asian Vegetable Resources 2: Edible Fruits. Jakarta (ID): Gramedia Main Library.
- 15 Verheij, EWM and RE Coronel., 1992. Plant Resources of South East Asia. 2: Edible Fruit and Nut. Prosea, Bogor.
- Victoria, H, 2010. Cultivation and Increase in Selling Value of Guava in Jogotirto Hamlet, Krasakan Village, Berbah District, Sleman Regency. Essay. Yogyakarta State University, Yogyakarta. Matters: 3-13.
- 6 Widowati. LR, Widati, S, U. Jaenudin, W. Hrtatik. 2004. Effect of organic fertilizer compost that is enriched with mineral and biological fertilizers on soil properties, nutrient uptake and organic vegetable production. Report of the Research Project for Agribusiness Development Program. Soil Research Center.
- Wijayanti, 2013. The Effect of Chicken Manure and Goat Manure on the Productivity of Chili Pepper (*12* *Psicum Frutescens* L.). Biology Education Study Program, Teacher Training and Education Faculty of Muhammadiyah University of Surakarta,
- Zakiah, M., TF Manurung and Wulandari Hospital. 2018. Chlorophyll Leaf Content in Four Tree Types at Sylva Indonesia Pe Arboretum. Tanjungpura University. Journal of Sustainable Forests (2018) Vol. 6 (1): 48 - 55. Faculty of Forestry, Tanjungpura University, Pontianak.

THE EFFECT OF PRUNING AND CHICKEN MANURE ON VEGETATIVE GROWTH OF HONEY DELI (*Syzygium aqueum* Burn F.) IN 9 MONTHS AGE

ORIGINALITY REPORT

10%

SIMILARITY INDEX

%

INTERNET SOURCES

10%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

A R Fahreza, N Aini. "Growth and fruit yield of sponge gourd (*Luffa acutangula* L.) as influenced by combination of plant spacing and chicken manure", IOP Conference Series: Earth and Environmental Science, 2023

Publication

2%

2

Nur Humairoh Arzad, Yohanis Tambing, Bahrudin Bahrudin. "THE EFFECT OF VARIOUS RATES OF COW MANURE APPLICATION ON GROWTH AND YIELD OF MUSTARD (*BRASSICA JUNCEA* L.)", AGROLAND: The Agricultural Sciences Journal, 2018

Publication

1%

3

S R Siswana, M Sembiring, H Hanum, Rosneli. "The effect of phosphate solubilizing microbes and chicken manure in increasing the P availability and growth of Green Beans (*L.*) on Andisol ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

1%

4

Nurzainah Ginting, Hasnudi, Yunilas. " Biogas slurry application on : A microbial transportation to improve plant quality ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

1 %

5

M F Ramadhani, A E Prihatiningrum, M. Abror. "Optimization of Mustard Pakcoy Growth: The Influence of Media Composition and Coconut Water Concentration", IOP Conference Series: Earth and Environmental Science, 2023

Publication

1 %

6

Perkasa, Achmad Yozar, Endang Gunawan, Sangrani Annisa Dewi, and Ulya Zulfa. "The Testing of Chicken Manure Fertilizer Doses to Plant Physiology Components and Bioactive Compound of Dewa Leaf", Procedia Environmental Sciences, 2016.

Publication

1 %

7

T E N Siagian, E R Sasmita, E B Irawati. "The Growth and Yield Responses of Shallot (*Allium Ascalonicum* L.) to Plant Spacing and Tuber Cutting by NFT Hydroponic", IOP Conference Series: Earth and Environmental Science, 2022

Publication

1 %

8

"Book of Abstracts International Conference on Agriculture, Environment, and Food

1 %

9

Alridiwersah, E M Harahap, E N Akoeb, H Hanum. "Growth and production of new superior rice varieties in the shade intensity", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

10

E B Irawati, E R Sasmita, A Suryawati. "Application of chitosan for vegetative growth of kemiri sunan plant in marginal land", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

11

Musfal. "The agricultural sustainability in rainfed land through the application of large super technology packages", IOP Conference Series: Earth and Environmental Science, 2023

Publication

<1 %

12

Aulia Hanifah Qomar. "AN ANALYSIS OF THE THIRD SEMESTER STUDENTS' ABILITY IN DEVELOPING NARRATIVE TEXT OF ENGLISH EDUCATION STUDY PROGRAM IN MUHAMMADIYAH UNIVERSITY OF METRO", PREMISE JOURNAL:ISSN online: 2442-482x, ISSN printed: 2089-3345, 2016

Publication

<1 %

13

L M Mooy, A Hasan, R Onsili. " Growth and yield of Tomato as influenced by the combination of liquid organic fertilizer concentration and branch pruning ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

14

Luciana Delgado-Benarroch. "The mutants compacta ähnlich, Nitida and Grandiflora define developmental compartments and a compensation mechanism in floral development in *Antirrhinum majus*", Journal of Plant Research, 05/02/2009

Publication

<1 %

15

Euis H. Hakim, Sjamsul A. Achmad, Lia D. Juliawaty, Lukman Makmur et al. "Prenylated flavonoids and related compounds of the Indonesian *Artocarpus* (Moraceae)", Journal of Natural Medicines, 2006

Publication

<1 %

16

B N Fitriatin, S Febriani, A Yuniarti. "Application of biofertilizers to increase upland rice growth, soil nitrogen and fertilizer use efficiency", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

17

Behera, S.K.. "Evaluation of plant performance of *Jatropha curcas* L. under

<1 %

different agro-practices for optimizing biomass A case study", Biomass and Bioenergy, 201001

Publication

18

D U Siswanti, N Umah. "Effect of Biofertilizer and Salinity on Growth and Chlorophyll Content of Amaranthus tricolor L", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

19

"Editorial Board", Wiley, 2005

Publication

<1 %

20

Biruk Masrie, Tesfaye Girma. "Effect of stem pruning and staking time on growth and yield of tomato (Lycopersicon esculentum Mill. Var. Mersa) at Mersa, North Wollo, Ethiopia", Vegetos, 2022

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off